

Examination of high-resolution feed intake data of grower finisher pigs confronted with typical short-term disturbances in stable routine

PETER LOIBL^{1*}, WILHELM WINDISCH¹, WOLFGANG PREIßINGER²

¹TUM School of Live Science Weihenstephan, Chair for Animal Nutrition, Technical University of Munich, Freising-Weihenstephan, Germany

²Institute for Animal Nutrition, Bavarian State Research Center for Agriculture, Grub, Germany

*Corresponding author: peter.loibl@wzw.tum.de

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic Supplementary Material (ESM)

Figure S1. Average daily feed intake per animal and fattening period; tick marks indicate standard deviations

Figure S2. Average daily feed intake per feeder visit, animal and fattening period; tick marks indicate standard deviations

Figure S3. Average daily number of visits to feeder per animal and fattening period; tick marks indicate standard deviations

Figure S4. Average daily feed intake of the single feeding action with highest feed intake per animal and fattening period; tick marks indicate standard deviations




Figure S1. Average daily feed intake per animal and fattening period; tick marks indicate standard deviations




Figure S2. Average daily feed intake per feeder visit, animal and fattening period; tick marks indicate standard deviations




Figure S3. Average daily number of visits to feeder per animal and fattening period; tick marks indicate standard deviations




Figure S4. Average daily feed intake of the single feeding action with highest feed intake per animal and fattening period; tick marks indicate standard deviations