

<https://doi.org/10.17221/116/2025-CJAS>

Mitigating methane in dairy cattle: Integrated strategies and the evolving role of precision livestock farming

EVA MIXTAJOVÁ¹, JOANA NERY², RADOVAN KASARDA¹, MUZAFFER DENLI⁴,
ACHILLE SCHIAVONE², ALKAN ÇAĞLI⁴, JOSÉ FRANCISCO PÉREZ⁵,
HASAN HÜSEYİN İPÇAK⁴, JOSÉ LUIS REPETTO⁵, STANISLAVA DROTÁROVÁ³,
CECILIA CAJARVILLE^{5*}

¹*Faculty of Agrobiological and Food Resources, Project Management Office, Slovak University of Agriculture in Nitra, Nitra, Slovak Republic*

²*Department of Veterinary Sciences, University of Turin, Grugliasco, TO, Italy*

³*Institute of Nutrition and Genomics, Slovak University of Agriculture in Nitra, Nitra, Slovak Republic*

⁴*Department of Animal Science, Faculty of Agriculture, Dicle University, Diyarbakir, Türkiye*

⁵*Department of Animal and Food Science, Faculty of Veterinary Medicine, Autonomous University of Barcelona – UAB, Barcelona, Spain*

*Corresponding author: ccajjarville@gmail.com

Citation: Mixtajová E., Nery J., Kasarda R., Denli M., Schiavone A., Çağlı A., Pérez J.F., İpçak H.H., Repetto J.L., Drotárová S., Cajarville C. (2026): Mitigating methane in dairy cattle: Integrated strategies and the evolving role of precision livestock farming. *Czech J. Anim. Sci.*, 71: 41–58.

Abstract: Modern dairy farming faces the dual challenge of meeting global food demands while mitigating its environmental impact, particularly greenhouse gas (GHG) emissions, such as methane (CH₄), a potent contributor to climate change. This review explores the role of Precision Livestock Farming (PLF) technologies in monitoring and reducing CH₄ emissions from dairy cattle. We evaluate state-of-the-art methods, including direct monitoring (e.g. respiratory chambers, GreenFeed systems) and indirect approaches (e.g. infrared milk spectroscopy, AI-driven analytics), alongside mitigation strategies such as nutritional optimisation, genetic selection, and ruminal additives. PLF emerges as a transformative tool, integrating real-time data on animal health, feed efficiency, and environmental conditions to optimise management practices and reduce emissions per unit of milk produced. By synthesising current research, we highlight the potential of PLF to reconcile productivity with sustainability, offering scalable solutions for the dairy sector. Critical gaps in real-time CH₄ monitoring and farm-level implementation are identified, underscoring the need for further innovation. This review provides a roadmap for aligning dairy production with global climate goals while ensuring food security for the growing population.

Keywords: emission monitoring; mitigation strategies; rumen fermentation; ruminant emissions; sustainability

Supported by the GLISTEN Project (Green Livestock Farming through Smart Technology in HEIs), 2023-1-TR01-KA220-HED-000157708, Erasmus +, EU.

© The authors. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0).

GREENHOUSE GAS EMISSIONS IN DAIRY CATTLE PRODUCTION: A GLOBAL PERSPECTIVE

The global human population is projected to reach 10 billion by 2050, creating unprecedented demands for food production that are intensifying pressure on agricultural systems and natural resources (von Soosten et al. 2020). As livestock production expands to meet growing protein needs, its environmental impacts have become increasingly urgent, particularly greenhouse gas (GHG) emissions from ruminant systems. Within this landscape, dairy farming represents a significant but addressable contributor to climate change through multiple pathways, including feed production, manure management, and most notably, enteric methane (CH₄) formation (Rotz 2018). The methane exceptional warming potential, 84 times greater than that of carbon dioxide (CO₂) over 20 years (Myhre et al. 2013), and its substantial role in near-term climate change have made it a priority for global action, as evidenced by the Global Methane Pledge's commitment by over 160 countries to reduce anthropogenic CH₄ emissions by 30% before 2030 (Global Methane Pledge 2021). Although the

top CO₂-emitting countries do not fully overlap with the leading methane (CH₄) emitters from livestock (Figure 1), methane remains a critical climate concern, particularly in regions like the EU, where enteric fermentation from cattle alone accounts for 38% of total anthropogenic methane emissions (European Environment Agency & European Commission 2025). This underscores the outsized role of livestock in climate change and emphasises the need for targeted mitigation strategies. This is particularly critical because cattle (dairy and non-dairy) production systems contribute to over 70% of total livestock-related CH₄ emissions (Figure 2), with the primary source being enteric fermentation, the microbial breakdown of organic matter in the rumen and hindgut (Wattiaux et al. 2019). Notably, this process is a major emissions driver in both dairy and non-dairy systems (Figure 3), emphasising the broad relevance of mitigation efforts across ruminant production.

This review examines the current understanding of CH₄ emissions from dairy production, evaluates established and emerging mitigation strategies, and explores how Precision Livestock Farming (PLF) technologies can enhance emission reduction efforts. The principles discussed extend beyond dairy cattle to other ruminant systems, including beef production, sheep, and goats, making these insights broadly relevant for sustainable livestock agriculture. By synthesising research across these areas, we aim to identify practical, scalable solutions that can help align animal protein production with climate stabilisation goals while maintaining food security in an era of global population growth.




Figure 1. Worldwide top 10 emitters in 2022 for total CH₄ (A) and Livestock (B)

Adapted from WRI (Friedrich et al. 2023)




Figure 2. Worldwide total CH₄ emissions by livestock species and categories estimated by FAO for the year 2022 (data from FAOSTAT 2025)

<https://doi.org/10.17221/116/2025-CJAS>




Figure 3. Total methane emissions and enteric-origin methane emissions estimated by FAO for 2022 in dairy and non-dairy cattle (FAOSTAT 2025)

ENTERIC METHANOGENESIS IN RUMINANTS: THE ENVIRONMENTAL–NUTRITIONAL PARADOX

The unique ruminant digestion system, which upgrades low-value feedstuffs into high-quality protein, is intrinsically linked to enteric methanogenesis. This microbial process, essential for efficient fibre degradation, results in the inevitable production of the potent greenhouse gas CH₄. Consequently, a fundamental tension exists between using ruminants to achieve global food security and mitigating their impact on climate change.

The rumen is a symbiotic bioreactor where microbial fermentation breaks down ingested feed, primarily converting carbohydrates into volatile fatty acids (VFAs), the ruminant's primary energy source, and microbial protein, which becomes the animal's primary source of amino acids. This anaerobic process is crucial for the ruminant's ability to upcycle low-quality forage.

A critical consequence of this fermentation is the production of metabolic H₂ (hydrogen). The type of volatile fatty acid (VFA) produced in the rumen directly influences hydrogen (H₂) dynamics, which in turn governs methanogenesis. Acetate and butyrate production belongs among net generators of H₂, whereas propionate synthesis acts as an H₂ sink. Since H₂ accumulation is toxic to many fermentative microbes, the rumen ecosystem depends on methanogenic archaea to maintain metabolic balance. These archaea remove the excess H₂ by using it to reduce CO₂ to CH₄, making methanogenesis an indispensable process for sustaining the overall fermentation efficiency (Hungate 1966). The balance of VFA production

directly determines the theoretical potential for methane emission, making it a key target for nutritional mitigation strategies.

The archaea comprise 2–4% of rumen microbiota (Waters et al. 2025), of which an overwhelming majority (90%) are methanogens, predominantly *Methanobrevibacter* spp. (Danielsson et al. 2017). These microorganisms are fundamental to the rumen function by acting as the primary H₂ sink, using the H₂ produced during carbohydrate fermentation to reduce CO₂ to CH₄ via the reaction: CO₂ + 4H₂ → CH₄ + 2H₂O. As CH₄ is eructated and represents an energy loss of 2–12% for the animal, this process directly links ruminant production to greenhouse gas emissions, accounting for approximately 14.5% of all the GHG (Ghassemi Nejad et al. 2024).

The relationship between dietary fibre and CH₄ production in ruminants is fundamentally linked to shifts in rumen fermentation pathways (Janssen 2010; Bannink et al. 2011) and microbial ecology. High-fibre diets promote acetogenic fermentation, which generates H₂ (e.g. C₆H₁₂O₆ + 2H₂O → 2CH₃COOH + 2CO₂ + 4H₂), thereby increasing substrate availability for methanogens. Conversely, high-concentrate diets shift fermentation towards propionogenesis, which consumes H₂ (e.g. C₆H₁₂O₆ + 2H₂ → 2CH₃CH₂COOH + 2H₂O), thus competitively reducing its availability for CH₄ production.

Microbial communities adapt dynamically to these dietary changes: fibrolytic bacteria (e.g. *Fibrobacter succinogenes*) and other acetate producers increase in high-fibre conditions, while starch-rich diets select for amylolytic bacteria (e.g. *Streptococcus bovis*) and propionate-generating pathways (e.g. via succinate or acrylate routes) (Morgavi et al. 2010). Additionally, methanogen populations (e.g. *Methanobrevibacter*) decline when fibre is reduced, further suppressing CH₄ (Hook et al. 2010). However, leveraging this dietary principle for mitigation involves a critical trade-off, as low-fibre diets risk causing acidosis and impairing the overall rumen health, highlighting the importance of balanced nutritional strategies (Beauchemin et al. 2020).

Therefore, while ruminal methanogenesis is an intrinsic outcome of the symbiosis that allows ruminants to upgrade low-quality feed, it necessitates targeted nutritional strategies to balance production efficiency with climate objectives.

STRATEGIES FOR METHANE REDUCTION IN DAIRY CATTLE: PRINCIPLES AND EVIDENCE FROM RUMINANT RESEARCH

Despite the paradox mentioned above, the available evidence suggests that strategies to reduce CH₄ emissions in dairy cattle do not inherently conflict with improved productivity. From a theoretical standpoint, a reduction in CH₄ emission represents a lower loss of dietary energy through eructation. This saved energy could potentially be redirected toward production, meaning that strategies that enhance feed efficiency often lead to both improved productivity and reduced CH₄ yield (Lovendahl et al. 2018). The purpose of this item is to provide a general overview of possible strategies to reduce CH₄ emissions. For comprehensive reviews on the topic, we recommend Hristov et al. (2013), Beauchemin et al. (2020), or Almeida et al. (2021).

Nutritional strategies that enhance propionate production in the rumen

As explained before, the higher the proportion of propionate in the rumen, the lower the CH₄ emitted. In this sense, higher non-fibre carbohydrate (NFC) content in the diets promotes propionate synthesis, diverting H₂ away from CH₄ production (Janssen 2010; Bannink et al. 2011). However, it is well-known that high NFC diets pose a risk of acidosis, which is greater when diets are low in forage-derived fibre (NASEM 2021). In a meta-analysis of 73 studies, Herliatika et al. (2024) demonstrated that increasing dietary starch in cattle mitigates enteric CH₄ production by two ways of action: (i) altering rumen protozoa populations [key H₂ producers according to Lewis et al. (2020)] and (ii) shifting the rumen fermentation profile toward propionate at the expense of acetate.

While modulating the carbohydrate composition is one effective approach, modifying the dietary lipid content represents another major avenue for reducing enteric methane. Indeed, changes in the ration content and the addition of lipids to dairy cow diets, although changing the fermentation profile toward acetate formation (Szumacher-Strabel et al. 2002), have been shown to reduce CH₄ emissions, concurrently altering the fatty acid profile

in milk and demonstrating a positive correlation between these parameters (van Lingen et al. 2014).

It is generally accepted that providing highly digestible forages enhances efficient ruminal fermentation and reduces CH₄ emissions (Eugene et al. 2021). Forage maturity directly increases enteric CH₄ production through nutritional quality decline. As plants mature, the secondary cell wall development elevates lignin content and reduces digestibility, simultaneously depressing intake and energy availability while enhancing methanogenesis per feed unit (Thompson and Rowntree 2020). These structural changes simultaneously lower forage intake and energy availability, promoting greater methanogenesis per unit of ingested feed. Archimede et al. (2011) reported that cattle fed warm-season legumes produced the lowest CH₄ emissions per unit intake, followed by cool-season legumes and temperate (C3) grasses, with tropical (C4) grasses yielding the highest emissions. Wims et al. (2010) and Munoz et al. (2016), working with dairy cows grazing high or low pre-grazing herbage mass, found that low herbage mass swards improved grass quality and reduced CH₄ emissions per cow, per unit of milk, and kilogram of dry matter intake due to higher forage quality. In beef cattle, Dini et al. (2018) demonstrated that grazing on high-quality pastures reduced enteric CH₄ emissions by approximately 15% compared to low-quality pastures across seasons, aligning with observations in dairy herds. In contrast, their earlier work with grazing dairy cattle (Dini et al. 2012) found no significant differences in milk yield or CH₄ production between legume-rich and grass-rich pastures. The authors attributed this result to the high herbage allowance that enabled selective grazing by cows, highlighting the ability of grazing ruminants to improve the chemical composition of their diet through selective foraging. Additionally, certain plant components, such as tannins, can influence CH₄ emissions. In a grazing study with heifers, Alecrim et al. (2024) demonstrated that incorporating tannin-rich legumes into grasslands is a strategy for mitigation by efficiency. In fact, although these authors did not observe a reduction in the total CH₄ emitted per animal, this approach effectively lowered the carbon footprint of beef production by drastically reducing methane emissions per kilogram of weight gain.

Genetic enhancement of forage species through traditional breeding techniques or genetic modi-

<https://doi.org/10.17221/116/2025-CJAS>

fication can lead to important gains in nutritional quality (Capstaff and Miller 2018; Pereira et al. 2022). This field has grown rapidly in recent years, as advances in molecular sciences and biotechnology now provide powerful tools to uncover the molecular mechanisms behind key traits in forage species, facilitating breeding strategies and tools for creating and implementing enhanced genetic solutions (Bryan et al. 2024).

Ruminal additives

Significant research efforts in recent years have focused on identifying natural inhibitors and developing synthetic analogues to suppress CH₄ generation, with many products marketed as efficiency enhancers now claiming emission-reducing effects. Recent publications include reviews and guidelines for identifying and selecting emission-reducing additives (Durmic et al. 2025) and properly evaluating CH₄-mitigating additives in dairy cows (Hristov et al. 2025). Among emerging compounds, 3-nitrooxypropanol (3-NOP), which inhibits the enzyme methyl-coenzyme M reductase responsible for CH₄ formation, demonstrates a proven mitigation potential (studies report up to 30% CH₄ reduction), though some evidence suggests it may reduce production (Pupo et al. 2025). Since NDF reduces 3-NOP effectiveness, the compound works better in intensive systems where diets contain more concentrates than fibre (Kebreab et al. 2023; Xuan et al. 2024). Among natural additives, the most frequently studied have been tannins, saponins, essential oils, and seaweeds. Tannins showed the greatest effect (Almeida et al. 2021), while essential oils demonstrated positive effects when used in long-term studies (Belanche et al. 2020). The outcomes depend on phytochemical combinations, dosages, and basal diet composition (Beauchemin et al. 2020).

While some studies show no effects using essential oils (Benchaar and Hassanat 2025), others report a clear CH₄ reduction using blends of essential oils and tannins in *in vitro* (Rossi et al. 2022) and *in vivo* studies (Minutti et al. 2025). Seaweeds like *Asparagopsis taxiformis*, containing bromoform as the main active compound, have also been studied, showing favourable results (Glasson et al. 2022). Natural compounds remain promising, especially when combined with other strategies. Still,

challenges persist in identifying specific mitigants, optimising doses and synergies (Flachowsky and Lebziern 2012), and elucidating microbial mechanisms beyond the confirmed effects (Belanche et al. 2025).

Genetic selection of the animals for higher production efficiency

Methane emissions in dairy cows can be reduced through genetic selection, using either direct or indirect genetic traits (Wall et al. 2010). Direct genetic traits are individual and specifically related to CH₄ emissions, whereas indirect genetic traits are related to the herd performance. Because genetic improvements are both permanent and cumulative, they represent a particularly promising long-term strategy for mitigating greenhouse gas emissions from ruminant production systems.

Selecting programmes for higher feed efficiency has been demonstrated to be useful to choose to emit less CH₄, in both dairy and beef cattle (Lovendahl et al. 2018; Olijhoek et al. 2018; Dini et al. 2019; Jenkins et al. 2024). De Haas et al. (2011) demonstrated that genetic selection could reduce cattle CH₄ emissions by 11–26% over a decade, a strategy with the potential for synergistic effects when integrated with feed additives, pasture management, and precision farming technologies. A recent large-scale Dutch study (7 139 cows across 68 farms) using automated “sniffer” sensors for CH₄ measurements revealed moderate heritability of CH₄ concentration and weak genetic correlations with production traits, suggesting the breeding for CH₄ mitigation could complement productivity goals (van Breukelen et al. 2025). Selection programs targeting rumen microbiome-associated traits are also promising (Mizrahi et al. 2021). Wall et al. (2010) proposed genetic selection for fitness traits (health, fertility, longevity) as a CH₄ mitigation tool. In dairy systems, increasing the cow longevity from 3 to 3.5 lactations reduces heifer replacement numbers, shrinking the herd’s carbon footprint without compromising the production, a whole-herd genetic strategy rather than an individual-efficiency metric. This author estimated that this approach would lead to a decrease in CH₄ emission by 0.55% in milking cows and 18.87% in young stock, with a mean total decrease by 4.42% in the UK.

Herd management

We cannot overlook the numerous management measures that aim to optimise the production process, which, as a byproduct, also reduce CH₄ emissions. It is important to remember that low-producing animals or those in maintenance, though emitting less, continue to release CH₄. Furthermore, [Goopy et al. \(2020\)](#) found that feeding cattle below maintenance levels increases CH₄ production per unit of feed intake, suggesting a counterproductive effect on emission reduction strategies.

Reducing unproductive periods (e.g. rearing, dry cows), improving reproductive performance, enhancing herd health, and minimising stress are herd tools that directly contribute to lowering CH₄ emissions by improving feed-to-product conversion efficiency ([Pulina et al. 2017](#)).

For instance, in a study conducted on Scottish dairy farms, [Ferguson et al. \(2024\)](#) reported that improving reproductive efficiency and reducing disease incidence could decrease total GHG emissions by 5–12%, and this magnitude was dependent on the baseline management conditions. [Beauchemin et al. \(2020\)](#) reached similar conclusions after reviewing 50 years of studies on methanogenesis in ruminants. These authors, however, caution that it is also unclear whether a decrease in CH₄ production leads to a consistent and sustained improvement in animal performance over time, information that will be necessary for producers to adopt. This information will also require further research on animal health, reproduction, product quality, cost-benefit ratio, safety, and consumer acceptance. At the same time, precision feeding, the practice of tailoring diets to individual animals, can minimise nutrient waste and associated emissions, although its implementation remains challenging in extensive production systems ([Chase and Fortina 2023](#)).

Manure and slurry management

Greenhouse gas emissions from ruminant manure and slurry depend on their composition, storage conditions, treatments, and soil distribution methods. Moreover, the management of manure and slurry can be independent of the dairy production efficiency and can be applied despite the farming performance.

Storage conditions, such as temperature and duration, are critical, as demonstrated by [Cardenas et al. \(2021\)](#). These authors reported that temperature and storage duration are critically interlinked in controlling methane emissions. While the long storage in warm summers led to high emissions, the same long duration in cold winters resulted in negligible release. The key mechanism is a temperature threshold of ~14 °C. Storage below this point permanently suppresses the manure methane potential, making the long-term cold storage an effective mitigation strategy where the factor of time works to prevent, rather than to cause, emissions. Therefore, they suggest that the yearly temperature sum should be monitored because the winter temperature influences the onset of methane emissions from slurry storage during the warm seasons. This agrees with [Balde et al. \(2016\)](#), suggesting anticipating the slurry pit management in late summer or early autumn to mitigate CH₄ emission. Another strategy to decrease CH₄ emissions from slurry includes reducing the manure liquid-phase or slurry volume in sump pits during high-temperature seasons ([Balde et al. 2016](#)). Higher CH₄ emissions were observed in covered slurry storage pits, probably due to an increased slurry temperature as compared to uncovered basins ([Vechi et al. 2023](#)).

Different treatments, physical, chemical and microbiological ones, can be applied to change the manure and slurry composition and consequently mitigate CH₄ emissions. Treatments aiming to decrease organic matter or methanogenesis in slurry and manure, such as anaerobic digestion, acidification, or composting, contribute to reducing the storage and field release of CH₄. Slurry chemical amendments can be used to decrease methanogen activity in slurry and significantly reduce CH₄ emissions during slurry storage. For instance, [Sommer et al. \(2017\)](#) proposed to reduce methane emissions by 68% using the sulphuric acid treatment and decreasing slurry pH to 5.5. Finally, the distribution methods of manure and slurry can lead to a variation in GHG emissions from the soil. However, according to [Sherman et al. \(2021\)](#), unlike other GHGs, CH₄ emissions are not influenced by manure and slurry distribution techniques but vary with soil moisture and temperature. Precision agriculture methods can be used to optimise the distribution of manure and slurry in the fields according to soil organic matter and fertility.

<https://doi.org/10.17221/116/2025-CJAS>

Manure management practices also play a crucial role: strategies such as frequent manure removal from pens, solid-liquid separation, slurry acidification, and anaerobic digestion can significantly mitigate CH₄ emissions. For example, acidifying cattle slurry can reduce CH₄ emissions by up to 60% (Petersen et al. 2013). The authors suggest this practice is best suited for intensive production systems, where they identify the greatest potential for reducing methane emissions. In addition to acidification, anaerobic digestion, depending on the conditions of humidity, temperature, and organic matter, allows biogas recovery and cuts emissions from storage by 40–80% depending on the system efficiency (Montes et al. 2013). These practices, while variable in applicability across systems, highlight that improvements in the herd and manure management represent effective, often complementary pathways to nutritional mitigation strategies.

It is known that feeding practices influence the potential CH₄ emissions of manure. High-forage diets increase the manure organic matter content, leading to a potentially higher CH₄ emission compared to high-concentrate systems. In fact, the concentration of non-digestible neutral detergent fibre (NDF) by *in situ* incubation has been proposed as one method to estimate CH₄ emissions from dairy cow manure (Huhtanen et al. 2021). Furthermore, evidence from tropical forage grasses links the low nitrogen content in forage to reduced CH₄ emissions from manure, an effect attributed to the suppression of methanogens under low N availability (Leitner et al. 2021).

Several research studies previously reviewed by Owen and Silver (2015) demonstrate that methane emissions from slurry and manure are underestimated by the most widely used estimation systems, such as the IPCC used to monitor national methane emissions. Therefore, on-site monitoring of methane emissions is fundamental to assessing the efficacy of nutritional, animal management, and manure management to mitigate emissions of methane. Monitoring sensors applied to flux chambers and wind tunnels can be used to measure CH₄ emissions from barns, manure collection platforms, slurry storage basins, or in the field (Ding et al. 2016; Figure 4). Moreover, considering that methane emissions are highly influenced by slurry physicochemical conditions, monitoring slurry temperature and pH in slurry basins could be a cost-effective measure to program slurry

treatments throughout the warm season, when the environmental conditions are favourable for methanogen activity.

METHANE MONITORING TECHNOLOGIES: FROM ANIMAL TO FARM LEVELS

Various methods exist to monitor and quantify CH₄ emissions in dairy cows, each with distinct applications, advantages, and limitations. However, selecting the appropriate methodology requires more than only choosing a measurement technique, but also it demands alignment with the study objectives. The optimal approach will differ depending on whether the data will be used for farm management decisions, feed evaluation, additive efficacy testing, or genetic selection. The intended application dictates the most suitable methodology. The first consideration should be whether direct animal measurements are required. There exist multiple *in vitro* methods that eliminate animal use, avoiding welfare concerns while reducing costs compared to *in vivo* approaches.

These techniques are particularly useful for preliminary assessments, such as initial feedstuff evaluations, additive screenings, or dose-response trials. *In vitro* approaches frequently allow deeper mechanistic studies, as researchers can fully control experimental conditions, including fermentation, substrate use, and digestive transit. However,




Figure 4. Experimental flux chamber and wind tunnel (extracted from Parker et al. 2013)


in vivo validation remains essential for conclusive results in later stages of research or product development. However, *in vivo* validation remains essential for conclusive results, despite the well-documented challenges of scaling reference methods like respiration chambers and the SF₆ tracer technique for practical, continuous on-farm use (Tedeschi et al. 2022).

For *in vivo* studies, methodologies can be broadly categorised as direct or indirect quantification approaches (Huhtanen et al. 2015). Direct quantification approaches measure the physical gas emissions from the animal, typically through breath analysis or chamber confinement. In contrast, indirect approaches estimate emissions through models based on proxies like milk mid-infrared (MIR) spectra or animal performance data. Three direct methods have been predominantly recognised as the most appropriate for regulatory and scientific purposes: the respiration chambers, the sulphur hexafluoride tracer technique, and the GreenFeed system (Thompson and Rowntree 2020; Hristov et al. 2025).

Respiration chambers, historically used in ruminant experiments for energy metabolism and nutrient utilisation studies, are regarded as the gold standard for quantifying emissions (Huhtanen et al. 2015; Ghassemi Nejad et al. 2024). The method involves housing animals in sealed chambers (Figure 5) where gas concentrations (e.g. CH₄, CO₂) are precisely monitored by comparing inlet and outlet air compositions. Accurate measurements require the proper instrument calibration and account for differences between incoming and outgoing gas concentrations, because air extraction rates and chamber piping systems can influence the precision of the results (Arceo-Castillo et al. 2019). The main limitation is related to the artificial environment and restricted movement imposed on animals, which may not accurately represent natural production settings. This raises concerns about biological validity, especially for extrapolations in grazing conditions. Furthermore, the necessity for animal acclimation and training introduces additional variables that could compromise data extrapolation to real-world scenarios. Consequently,



(A)



(C)



(B)

Figure 5. Chambers of the ILVO Ruminant Respiration Facility

(A) Four of the six chambers; (B) Inside view of a chamber made of polypropylene mounted on a stainless-steel frame; (C) Cow tied in the chamber (extracted from: De Campeneere and Peiren 2014)

<https://doi.org/10.17221/116/2025-CJAS>

while respiration chambers provide highly precise data under controlled conditions, the artificial environment introduces a degree of systematic uncertainty when these measurements are extrapolated to predict emissions in commercial, free-moving herds. This limitation necessitates a cautious interpretation of absolute emission values for on-farm inventory purposes.

The sulphur hexafluoride (SF_6) tracer technique (Johnson et al. 1994) quantifies enteric CH_4 emissions by placing a slow-release capsule of SF_6 in the rumen, which disperses at a known rate. A sampling device worn by the animal continuously draws air from the muzzle into a canister over 3–6 days, capturing both CH_4 and SF_6 (Figure 6). By analysing the ratio of these gases in the collected sample and referencing the predetermined SF_6 release rate, total CH_4 production can be accurately estimated. This portable, low-cost system is suitable for both grazing and confined animals, offering practical advantages in diverse settings. It has been adopted in several regions for its non-invasive nature and reliability in field studies (Della Rosa et al. 2021). However, this method relies on labour-intensive custom-built sampling systems to assemble from heterogeneous components. The equipment requires continuous monitoring to prevent animal dislodgment, further increasing the operational de-

mands of data collection and analysis. Additionally, it is less reliable for assessing daily emission patterns and involves the use of trained animals. Therefore, while the SF_6 technique provides valuable field-based data, the cumulative effect of these operational demands can introduce a non-trivial degree of measurement uncertainty. This underscores a limitation for its application in generating the highly consistent, granular data required to formulate specific, on-farm mitigation strategies.

The GreenFeed system completes the trio of established measurement approaches. This automated technology, first developed by Zimmerman (1993) and later upgraded in 2012 (Zimmerman and Zimmerman 2012), quantifies enteric CH_4 emissions during ruminant feeding through controlled airflow sampling. The system captures real-time data, typically requiring ~50 samples over two weeks for reliable estimates. Unlike respiration chambers, it allows animals to move freely, but measurements are restricted to trained animals visiting the unit (Hammond et al. 2016). A key limitation is its reliance on concentrate-based rewards to attract animals, which means emissions are sampled only during supplemental feeding and not during forage-only diets. The system's hardware and data processing are vendor-specific (C-Lock Inc., Rapid City, South Dakota, USA), which could



Figure 6. Using the SF_6 tracer method in the IPAV (Veterinary Faculty, UdelaR, Uruguay, images from the authors)

(A) How the system works: a capsule placed in the rumen releases SF_6 at a known rate (in green), and a sampling device in the muzzle of the animal continuously draws exhaled air from the muzzle of the animal into a vacuum canister for 3–6 days (in yellow); (B) Detail of the vacuum canister and the sampler; (C) Milking cow on pasture during a measurement period

limit customisation and raw data access if necessary. While studies have demonstrated that the GreenFeed system provides accurate estimates of emission rates (McGinn et al. 2021), this accuracy is confined to the specific conditions of measurement. The combination of voluntary animal interaction, potential site-specific interference, and the sampling of emissions primarily during supplemental feeding bouts introduces a significant uncertainty regarding the representativeness of these measurements for the total daily methane production of the herd under natural conditions.

A comprehensive review by Della Rosa et al. (2021) analysing approximately 400 publications from 1995–2018 found that respiration chambers were the predominant technique (55% of studies), followed by the SF₆ tracer method (38%), with GreenFeed systems representing just 7% of studies. However, these proportions will likely shift in future analyses given the increasing adoption of GreenFeed systems by the global research community, with installations continuing to grow due to their advantages for long-term CH₄ monitoring in production environments. This historical reliance on methods with significant trade-offs between precision (chambers) and real-world applicability (SF₆) underscores a key uncertainty in the existing literature: the challenge of generating datasets that are both highly precise and directly representative of commercial farm conditions.

Portable analysers like “sniffers” have been developed to measure emissions in real-time at the feed bunk (Garnsworthy et al. 2012). While practical for on-farm use, these systems are prone to significant inaccuracies. Unpredictable cow head movements, variations in the feed trough design, and inconsistent sampling point placement can all introduce substantial measurement uncertainty, raising questions about their reliability for precise quantification.

The field is rapidly evolving, with new devices in permanent development. For instance, the ZELP device, a field-deployable system using near-infrared sensors on a face mask to analyse exhaled breath, demonstrated a strong positive correlation ($r = 0.912$, $P < 0.05$) with gold-standard respiration chamber measurements (Coetzee and Bica 2025). While promising, its practical implementation depends on the consistent wearing of the mask by the animal, a variable that must be accounted for. Similarly, hand-held laser detectors are

another portable method, noted for being relatively inexpensive and providing immediate results (Roessler et al. 2018; Kang et al. 2022). However, their measurements are highly sensitive to the operator’s distance and angle to the animal, introducing another layer of operational uncertainty.

In summary, while these emerging portable technologies offer compelling advantages for farm-level deployment, their current limitations – chiefly their susceptibility to environmental and behavioural variables – mean that rigorous on-farm validation and a clear understanding of their associated measurement errors are prerequisites for their reliable use in research and mitigation strategy assessment.

The use of artificial intelligence (AI) represents a promising indirect method for determining CH₄ emissions. Recent advances have demonstrated its potential, with estimations being improved by techniques such as support vector regression, random forests, and artificial neural networks that can explore complex, non-linear relationships in data (Ghassemi Nejad et al. 2024). However, the robustness of these models is heavily contingent on the quality, volume, and representativeness of the input data used for training. A key limitation and a source of uncertainty are the “black box” nature of some complex models, which can make it difficult to interpret the underlying physiological drivers of the predicted emissions. Furthermore, models trained on data from one production system or breed may not generalise well to others, creating a significant challenge for the broad farm-level implementation without extensive, and costly, re-validation.

Another indirect method for monitoring CH₄ emissions in dairy cattle integrates mid-infrared spectroscopy and milk fatty acid analysis (Engelke et al. 2018) with rumen volatile fatty acid profiles and microbiome characterisation (Bilton et al. 2025). This combined approach facilitates large-scale CH₄ quantification and could optimise breeding strategies for CH₄ reduction through genetic selection. The established relationships between milk fatty acid profiles and CH₄ emission levels (van Lingen et al. 2014) support the validity of these proxy measurements. However, a significant source of uncertainty stems from the confounding influence of other factors on these proxies. Key variables such as diet composition, energy balance, and health status can directly alter milk fatty acid profiles and rumen microbial communities inde-

<https://doi.org/10.17221/116/2025-CJAS>

pendently of changes in methanogenesis. This can lead to situations where a shift in a proxy variable is misinterpreted as a change in CH₄ production, thereby compromising the accuracy of animal phenotyping for genetic selection programs.

In conclusion, the selection of a CH₄ monitoring method is not a search for a universal optimum, but a strategic decision that must align the inherent strengths and limitations of the method with the specific objectives and constraints of the intended application, from fundamental research to on-farm management.

PRECISION LIVESTOCK FARMING AS A STRATEGIC TOOL FOR METHANE MITIGATION IN DAIRY SYSTEMS

The application of process engineering principles to livestock production management represents a transformative modern approach, systematically capturing the key operational parameters and efficiency metrics that define PLF (Tuytens et al. 2022). These technologies enable farmers to enhance operational efficiency while simultaneously addressing environmental concerns, safeguarding livelihoods, and improving animal health and welfare. Amid growing economic and environmental pressures in livestock production, PLF systems provide farmers with advanced tools to monitor animals more effectively, optimise welfare, and extend productive lifespans through smart technological solutions (Norton et al. 2019).

The most common PLF technologies employed on farms include the use of sensors as rumen monitoring boluses (Tuytens et al. 2022), oestrus detectors, in-line milk analysers, and automated lameness sensors (Ferguson et al. 2024), alongside environmental monitors (e.g. thermometers, acoustic sensors, cameras) tracking housing conditions. These on-farm tools enable real-time tracking of animal physiology, health, and environmental factors, allowing the recording of large amounts of data and information about animals and their management to optimise management decisions (Tuytens et al. 2022). The data can be used to adjust farm conditions and management in a short period, for example by adjusting feed rations and preventing metabolic diseases (rumen boluses), changing ventilation systems and adjusting ventilation slats (thermometers, airflow sensors), milking

(robotic systems) and reproductive management (animal movement and temperature sensors) (Norton et al. 2019).

While PLF systems provide measurable improvements to livestock management, their implementation requires the careful risk assessment. Concerns include direct animal welfare impacts, such as health complications from sensor malfunctions or prolonged device exposure, and operational challenges like erroneous decisions caused by data misinterpretation. A significant secondary effect is the potential decline in husbandry skills and reduced human oversight as over-dependence on automated systems diminishes hands-on animal care practices. Based on this, the effective implementation of PLF systems requires continuous attention by the livestock breeder and careful selection of suitable PLF technologies (Tuytens et al. 2022).

The PLF technologies were primarily developed for reproductive and health monitoring, but they also offer a strategic potential for CH₄ mitigation in dairy systems due to their core features: real-time monitoring, automation, and data analytics. A range of PLF tools is currently utilised, including sensors that monitor behavioural and physiological parameters (activity, rumination, temperature, body condition), automated feeders for precision nutrition, and AI-based models capable of interpreting complex data streams for timely decision-making (Jacobs and Siegford 2012; Spoliansky et al. 2016; Strapak et al. 2021). Among these, rumen boluses have proven to be particularly valuable for the continuous monitoring of ruminal pH, temperature, and activity levels, providing critical insights into the animal's metabolic status (Studer et al. 2023; Vladimirov et al. 2023; Vakulya et al. 2024; Weir et al. 2024). These devices enable early detection of subacute ruminal acidosis and related disorders, which, if left unmanaged, can result in significant productivity losses and increased environmental impact (Voulgarakis et al. 2024; Pfrombeck et al. 2025). Furthermore, insights from bolus data support dietary adjustments that improve rumen stability and reduce CH₄ emissions associated with inefficient fermentation. For example, deviations from the optimal pH range have been associated with milk production losses of 6.8% (low pH) and 14.08% (high pH) (Hanusovsky et al. 2018) and therefore, the use of rumen sensors can help to lower the CH₄ emis-

sion intensity (i.e. CH₄ per kg of milk, fat or protein produced). Boluses have also been used to evaluate drinking behaviour, revealing that high ambient temperatures increase the water intake frequency, particularly after feeding and milking (Hanusovsky et al. 2017), which can adjust strategies for managing the heat stress that reduces feed efficiency. Although slight sensor drift and inter-system variability are limitations (Studer et al. 2023; Weir et al. 2024), these technologies remain reliable for long-term data collection.

In parallel, PLF tools such as motion and rumination sensors enhance reproductive efficiency by accurately detecting oestrus and calving, thereby reducing open days and improving productivity (Stevenson et al. 2014; Mayo et al. 2019; Strapak et al. 2021). Health monitoring systems, including automated body condition scoring (Fischer et al. 2015; Spoliansky et al. 2016) and infrared thermography for early mastitis and lameness detection (Zaninelli et al. 2018; Korelidou et al. 2024), further support the herd management efficiency. Collectively, although CH₄ mitigation is not the primary function of these systems, their application in optimising health, nutrition, and productivity aligns with sustainability goals by reducing CH₄ emissions per unit of milk produced. Ferguson et al. (2024) modelled the potential emission reductions using the PLF technologies available to farms and concluded that sensors used to improve fertility had the largest reduction potential to reduce the emissions expressed as kg CO₂ equivalent. In line with this, McNicol et al. (2024) demonstrated that the adoption of PLF technologies, specifically automatic weighing platforms, oestrus detection sensors, and health monitoring sensors, led to significant reductions in both total emissions (kg CO₂e) and product emissions (kg CO₂e/kg deadweight) in Scottish beef systems. The greatest reduction in total emissions was observed with weighing platforms in grazing systems (6.8%) and health sensors in housed systems (6.1%). In terms of product emissions, health sensors achieved the largest decrease (12% in housed systems and 10.5% in grazing systems). These findings suggest that PLF technologies can serve as an effective greenhouse gas mitigation strategy in dairy production systems.

To critically assess the role of PLF in methane mitigation, it is essential to distinguish between its current indirect contributions and its future

potential for direct management. Currently, PLF technologies contribute to methane emission mitigation primarily through indirect pathways. For example, by improving animal health and fertility, shortening time to slaughter, and increasing feed efficiency, all of which reduce CH₄ emissions per unit of product (Papakonstantinou et al. 2024).

In contrast, the development of PLF for direct methane measurement and mitigation is an ongoing research frontier. Direct measurement of CH₄ at the individual animal level on farms is possible using several new PLF tools, but these approaches still require further validation, cost reduction, and integration before they can deliver the routine emission mitigation on farms. Established direct methods (breathing chambers, SF₆ tracers) remain the reference standards but they are not practical for continuous monitoring on farms (Tedeschi et al. 2022). Practical systems on farms that provide near real-time measurements include automated sampling devices at head/feeding stations (e.g. GreenFeed) that capture exhaled air during voluntary visits and estimate individual enteric CH₄ fluxes, handheld/vehicle-mounted laser methane detectors (LMD/TDLAS) that detect breath vapours or emissions from barns, and fixed or mobile cavity ring-mode/dual-comb/open-path spectrometers that, when combined with wind/dispersion modelling, can quantify CH₄ fluxes in a barn or across the entire farm (McGinn et al. 2021). Each approach has its trade-offs in terms of accuracy, temporal resolution, animal coverage, and cost: GreenFeed and similar head-gate systems provide validated individual estimates, but sufficient animal visits and calibration are required to extrapolate to 24-hour emissions; LMD and mobile CRDS/TDLAS devices show promise for rapid screening measurements and farm-level flux estimation, but they require standardised protocols and quantification of uncertainty; open-path dual-comb systems and fixed CRDS systems show a potential for continuous whole-farm monitoring, but remain relatively expensive and require advanced data processing (Hristov et al. 2015). Crucially, however, these are not yet standard PLF tools for routine mitigation. Their widespread adoption faces significant hurdles, including the need for rigorous validation against gold-standard methods, the development of algorithms to convert spot measurements into reliable daily fluxes, substantial cost reduction, and farm trials dem-

<https://doi.org/10.17221/116/2025-CJAS>

onstrating that real-time data can be effectively translated into management actions that reduce methane (USDA 2024).

Therefore, while PLF could evolve into a direct methane monitoring tool, it currently functions most effectively as an indirect mitigation strategy. The critical challenge is to balance accuracy, animal welfare, and affordability to ensure the practical adoption. According to Kaur et al. (2023), future efforts should prioritise developing affordable, interoperable sensors and robust data integration. Ultimately, the transformation of animal agriculture toward greater sustainability will depend on merging these technological advancements with robust biological models to create closed-loop systems that not only monitor but also actively manage methane emissions.

CONCLUDING REMARKS

Over the past decades, numerous strategies to mitigate CH₄ emissions in dairy farming have been investigated. Both direct and indirect approaches, ranging from nutritional interventions and genetic selection to improved herd management, can enhance farm sustainability by optimising animal productivity and health, reducing CH₄ output. These approaches can enhance sustainability, but their effectiveness depends on precise implementation, a challenge where PLF could play a pivotal role. The PLF technologies offer tools to optimise these strategies on-farm, delivering direct benefits in productivity, while also enhancing farmers' decision-making and product traceability. Indirectly, PLF adoption can reduce the carbon footprint of livestock production and improve socio-economic outcomes. However, critical gaps remain, particularly in capitalising on PLF for real-time CH₄ monitoring, a key hurdle for targeted mitigation. For instance, automated CH₄ sensors paired with AI-driven analytics could provide actionable insights while reducing labour demands, but such solutions require further research and on-farm validation. To maximise the impact, future efforts must prioritise developing accessible, reliable PLF systems tailored to CH₄ measurement and farm-level decision support. Bridging this gap will be essential to translate mitigation strategies into practical, scalable solutions that align environmental goals with farm profitability and animal welfare.

Acknowledgement

We sincerely thank the authorities, staff, and students at the Autonomous University of Barcelona (UAB), Dicle University, the Slovak University of Agriculture in Nitra (SUA), and the University of Turin (UniTo) for their support in hosting the group meetings. This paper is the result of the project “Green Livestock Farming through Smart Technology in HEIs” (GLISTEN) (<https://glisten-project.eu/home>) funded by European Union Education and Youth Programs Center (Turkish National Agency); Grant No. 2023-1-TR01-KA220-HED-000157708.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Alecrim FB, Devincenzi T, Reyno R, Mederos A, Simon Zinno C, Mariotta J, Lattanzi FA, Nobrega GN, Santander D, Gere JJ, Irigoyen L, Ciganda VS. Addition of tannin-containing legumes to native grasslands: Effects on enteric methane emissions, nitrogen losses and animal performance of beef cattle. *Sustainability (Basel)*. 2024 Oct 16;16(20):9135.
- Almeida AK, Hegarty RS, Cowie A. Meta-analysis quantifying the potential of dietary additives and rumen modifiers for methane mitigation in ruminant production systems. *Anim Nutr*. 2021 Dec;7(4):1219-30.
- Arceo-Castillo JI, Montoya-Flores MD, Molina-Botero IC, Pineiro-Vazquez AT, Aguilar-Perez CF, Ayala-Burgos AJ, Ku-Vera JC. Effect of the volume of methane released into respiration chambers on full system methane recovery. *Anim Feed Sci Technol*. 2019 Jun;249:54-61.
- Archimede H, Eugene M, Magdeleine CM, Boval M, Martin C, Morgavi DP, Lecomte P, Doreau M. Comparison of methane production between C3 and C4 grasses and legumes. *Anim Feed Sci Technol*. 2011 Aug;166-167:59-64.
- Balde H, VanderZaag AC, Burt S, Evans L, Wagner-Riddle C, Desjardins RL, MacDonald JD. Measured versus modeled methane emissions from separated liquid dairy manure show large model underestimates. *Agric Ecosyst Environ*. 2016 Aug 16;230:261-70.
- Bannink A, van Schijndel MW, Dijkstra J. A model of enteric fermentation in dairy cows to estimate methane emission for the Dutch National Inventory Report using the IPCC

<https://doi.org/10.17221/116/2025-CJAS>

- Tier 3 approach. *Anim Feed Sci Technol.* 2011 Aug;166-167:603-18.
- Beauchemin KA, Ungerfeld EM, Eckard RJ, Wang M. Fifty years of research on rumen methanogenesis: Lessons learned and future challenges for mitigation. *Animal.* 2020 Mar;14(S1):s2-s16.
- Belanche A, Bannink A, Dijkstra J, Durmic Z, Garcia F, Santos FG, Huws S, Jeyanathan J, Lund P, Mackie RI, McAllister TA, Morgavi DP, Muetzel S, Pitta DW, Yanez-Ruiz DR, Ungerfeld EM. Feed additives for methane mitigation: A guideline to uncover the mode of action of anti-methanogenic feed additives for ruminants. *J Dairy Sci.* 2025 Jan;108(1):375-94.
- Belanche A, Newbold CJ, Morgavi DP, Bach A, Zweifel B, Yanez-Ruiz DR. A meta-analysis describing the effects of the essential oils blend Agolin Ruminant on performance, rumen fermentation and methane emissions in dairy cows. *Animals (Basel).* 2020 Apr 3;10(4):620.
- Benchaar C, Hassanat F. Diet supplementation with a mixture of essential oils: Effects on enteric methane emissions, apparent total-tract nutrient digestibility, nitrogen utilization, and lactational performance. *J Dairy Sci.* 2025 Apr;108(4):3560-72.
- Bilton TP, Hickey SM, Jonker A, McRae K, Hess MK, Perry BJ, Bryson B, Henry H, Bain W, Booker F, Veenvliet B, Peers-Adams J, Pile G, Waller E, Janssen PH, Tiplady KM, Reid P, Muetzel S, Agnew M, Dodds KG, McEwan JC, Rowe SJ. Milk fatty acids and rumen composition as proxy measures of enteric methane. *J Dairy Sci.* 2025 May;108(5):5125-44.
- Bryan G, Whitehayakul S, Roberts N. Nutritional enhancement of animal feed and forage crops via genetic modification. *J R Soc N Z.* 2024 Feb;55(2):327-42.
- Capstaff NM, Miller AJ. Improving the yield and nutritional quality of forage crops. *Front Plant Sci.* 2018 Apr 24;9:535.
- Cardenas A, Ammon C, Schumacher B, Stinner W, Herrmann C, Schneider M, Weinrich S, Fischer P, Amon T, Amon B. Methane emissions from the storage of liquid dairy manure: influences of season, temperature and storage duration. *Waste Manag.* 2021 Feb;121:393-402.
- Chase LE, Fortina R. Environmental and economic responses to precision feed management in dairy cattle diets. *Agriculture (Basel).* 2023 May;13(5):1032.
- Coetzee N, Bica R. Beyond respiration chambers: A field-deployable device for continuous methane emission measurement in cattle. In: *Book of Abstracts of the 76th Annual Meeting of the European Federation of Animal Science.* Wageningen (Netherlands): Wageningen Academic Publishers; 2025 Aug. p. 254.
- Danielsson R, Dicksved J, Sun L, Gonda H, Müller B, Schnurer A, Bertilsson J. Methane production in dairy cows correlates with rumen methanogenic and bacterial community structure. *Front Microbiol.* 2017 Feb 17;8:226.
- De Campeneere S, Peiren N. ILVO Ruminant Respiration Facility, Melle, Belgium. In: Pinares C, Waghorn G, editors. *Technical manual on respiration chamber designs* [Internet]. [place unknown]: Global Research Alliance on Agricultural Greenhouse Gases; 2014 Feb [cited 2025 Jan 23]. Available from: <https://globalresearchalliance.org/library/livestock-research-group-technical-manual-respiration-chamber-designs-february-2014/>.
- de Haas Y, Windig JJ, Calus MPL, Dijkstra J, de Haan M, Bannink A, Veerkamp RF. Genetic parameters for predicted methane production and potential for reducing enteric emissions through genomic selection. *J Dairy Sci.* 2011 Dec;94(12):6122-34.
- Della Rosa MM, Jonker A, Waghorn GC. A review of technical variations and protocols used to measure methane emissions from ruminants using respiration chambers, SF6 tracer technique and GreenFeed, to facilitate global integration of published data. *Anim Feed Sci Technol.* 2021 Jan;279:115018.
- Ding L, Cao W, Shi Z, Li B, Wang C, Zhang G, Kristensen S. Carbon dioxide and methane emissions from the scale model of open dairy lots. *J Air Waste Manag Assoc.* 2016 Jul;66(7):715-25.
- Dini Y, Briano C, Manetti M, Juliarena P, Picasso V, Gratton R, Astigarraga L. Methane emission and milk production of dairy cows grazing pastures rich in legumes or rich in grasses in Uruguay. *Animals (Basel).* 2012 Jun 8;2(2):288-300.
- Dini Y, Cajarville C, Gere J, Fernandez S, Fraga M, Pravia MI, Navajas E, Ciganda V. Association between residual feed intake and enteric methane emissions in Hereford steers. *Transl Anim Sci.* 2019 Jan 1;3(1):239-46.
- Dini Y, Gere JI, Cajarville C, Ciganda VS. Using highly nutritious pastures to mitigate enteric methane emissions from cattle grazing systems in South America. *Anim Prod Sci.* 2018 Dec;58(12):2329-34.
- Durmic Z, Duin EC, Bannink A, Belanche A, Carbone V, Carro MD, Crisemann M, Fievez V, Garcia F, Hristov AN, Joch M, Martinez-Fernandez G, Muetzel S, Ungerfeld EM, Wang M, Yanez-Ruiz DR. Feed additives for methane mitigation: Recommendations for identification and selection of bioactive compounds to develop anti-methanogenic feed additives. *J Dairy Sci.* 2025 Jan;108(1):302-21.
- Engelke SW, Das G, Derno M, Tuchscherer A, Berg W, Kuhla B, Metges CC. Milk fatty acids estimated by mid-infrared spectroscopy and milk yield can predict methane emissions in dairy cows. *Agron Sustain Dev.* 2018 Jun;38(3):29.

<https://doi.org/10.17221/116/2025-CJAS>

- Eugene M, Klumpp K, Sauvant D. Methane mitigating options with forages fed to ruminants. *Grass Forage Sci.* 2021 Sep;76(3):429-43.
- European Environment Agency & European Commission. Report: Annual European Union greenhouse gas inventory 1990–2023 and inventory report 2025 [Internet]. 2025 [cited 2026 Jan 20]. Available from: <https://www.eea.europa.eu/en/analysis/publications/annual-european-union-greenhouse-gas-inventory-2025>
- FAOSTAT – Food and Agriculture Organization of the United Nations. FAOSTAT: emissions from livestock [Internet]. Rome: FAO; 2025 [cited 2025 Jun 25]. Available from: <https://www.fao.org/faostat/en/#data/GLE>.
- Ferguson HJ, Bowen JM, McNicol LC, Bell J, Duthie CA, Dewhurst RJ. The impacts of precision livestock farming tools on the greenhouse gas emissions of an average Scottish dairy farm. *Front Sustain Food Syst.* 2024 May 22;8:1385672.
- Fischer A, Luginbühl T, Delattre L, Delouard JM, Faverdin P. Rear shape in 3 dimensions summarized by principal component analysis is a good predictor of body condition score in Holstein dairy cows. *J Dairy Sci.* 2015 Jul;98(7):4465-76.
- Flachowsky G, Lebzien P. Effects of phytogenic substances on rumen fermentation and methane emissions: A proposal for a research process. *Anim Feed Sci Technol.* 2012 Sep;176(1-4):70-7.
- Friedrich J, Ge M, Pickens A, Vigna L. This interactive chart shows changes in the world's top 10 emitters [Internet]. Washington (DC): World Resources Institute; 2023 [cited 2025 Jun 15]. Available from: <https://www.wri.org/insights/interactive-chart-shows-changes-worlds-top-10-emitters>.
- Garnsworthy PC, Craigon J, Hernandez-Medrano JH, Saunders N. Variation among individual dairy cows in methane measurements made on farm during milking. *J Dairy Sci.* 2012 Jun;95(6):3181-9.
- Ghassemi Nejad J, Ju MS, Jo JH, Oh KH, Lee YS, Lee SD, Lee HG. Advances in methane emission estimation in livestock: A review of data collection methods, model development and the role of AI technologies. *Animals (Basel).* 2024 Mar;14(3):435.
- Glasson CRK, Kinley RD, de Nys R, King N, Adams SL, Packer MA, Svenson J, Eason CT, Magnusson M. Benefits and risks of including the bromoform containing seaweed *Asparagopsis* in feed for the reduction of methane production from ruminants. *Algal Res.* 2022 Apr;64:102673.
- Global Methane Pledge. Global Methane Pledge: A commitment to reduce global methane emissions at least 30 percent from 2020 levels by 2030 [Internet]. 2021 [cited 2026 Jan 20]. Available from: <https://www.globalmethanepledge.org>.
- Goopy JP, Korir D, Pelster D, Ali AIM, Wassie SE, Schlecht E, Dickhoefer U, Merbold L, Butterbach-Bahl K. Severe below-maintenance feed intake increases methane yield from enteric fermentation in cattle. *Br J Nutr.* 2020 Jun 14;123(11):1239-46.
- Hammond KJ, Crompton LA, Bannink A, Dijkstra J, Yanez-Ruiz DR, O'Kiely P, Kebreab E, Eugene MA, Yu Z, Shingfield KJ, Schwarm A, Hristov AN, Reynolds CK. Review of current in vivo measurement techniques for quantifying enteric methane emission from ruminants. *Anim Feed Sci Technol.* 2016 Sep;219:13-30.
- Hanusovsky O, Biro D, Simko M, Galik B, Juracek M, Rolinac M, Balusikova E. The dynamic of the ruminal content pH change and its relationship to milk composition. *Acta Vet Brno.* 2018 Jun;87(2):119-26.
- Hanusovsky O, Biro D, Simko M, Galik B, Juracek M, Rolinac M, Herkel R. Drinking regime evaluation with continuous ruminal monitoring boluses. *Acta Fytotech Zootech.* 2017 Mar;20(1):1-5.
- Herliatika A, Widiawati Y, Jayanegara A, Harahap RP, Kusumaningrum DA, Shiddieqy MI, Adiati U. Meta-analysis of the relationship between dietary starch intake and enteric methane emissions in cattle from in vivo experiments. *J Adv Vet Anim Res.* 2024 Mar;11(1):212-30.
- Hook SE, Wright ADG, McBride BW. Methanogens: Methane producers of the rumen and mitigation strategies. *Archaea.* 2010 Dec 30;2010:945785.
- Hristov AN, Bannink A, Battelli M, Belanche A, Cajarville Sanz MC, Fernandez-Turren G, Garcia F, Jonker A, Kenny D, Lind V, Meale SJ, Meo D, Munoz C, Pacheco D, Peiren N, Ramin M, Rapetti L, Schwarm A, Stergiadis S, Theodoridou K, Ungerfeld EM, van Gastelen S, Yanez-Ruiz D, Waters SM, Lund P. Feed additives for methane mitigation: Recommendations for testing enteric methane-mitigating feed additives in ruminant studies. *J Dairy Sci.* 2025 Jan;108(1):322-55.
- Hristov AN, Oh J, Firkins JL, Dijkstra J, Kebreab E, Waghorn G, Makkar HP, Adesogan AT, Yang W, Lee C, Gerber PJ, Henderson B, Tricarico JM. Special topics – Mitigation of methane and nitrous oxide emissions from animal operations: I. A review of enteric methane mitigation options. *J Anim Sci.* 2013 Nov;91(11):5045-69.
- Hristov AN, Oh J, Giallongo F, Frederick T, Weeks H, Zimmerman PR, Harper MT, Hristova RA, Zimmerman RS, Branco AF. The use of an automated system (GreenFeed) to monitor enteric methane and carbon dioxide emissions from ruminant animals. *J Vis Exp.* 2015 Sep 7;(103):52904.
- Huhtanen P, Cabezas-Garcia EH, Utsumi S, Zimmerman S. Comparison of methods to determine methane emissions from dairy cows in farm conditions. *J Dairy Sci.* 2015 May;98(5):3394-409.

<https://doi.org/10.17221/116/2025-CJAS>

- Huhtanen P, Krizsan SJ, Ramin M. A meta-analysis of faecal output and nutrient composition, and potential methane emission from manure of dairy cows. *Anim Feed Sci Technol.* 2021 Dec;(282):115120.
- Hungate RE. *The rumen and its microbes.* New York: Academic Press; 1966.
- Jacobs JA, Siegford JM. Invited review: The impact of automatic milking systems on dairy cow management, behavior, health, and welfare. *J Dairy Sci.* 2012 May;95(5):2227-47.
- Janssen PH. Influence of hydrogen on rumen methane formation and fermentation balances through microbial growth kinetics and fermentation thermodynamics. *Anim Feed Sci Technol.* 2010 Aug;160(1-2):1-22.
- Jenkins B, Herold L, de Mendonca M, Loughnan H, Willcocks J, David T, Avis K. Breeding for reduced methane emissions in livestock. Edinburgh: ClimateXChange Publications; 2024. 64 p.
- Johnson K, Huyler M, Westberg H, Lamb B, Zimmerman P. Measurement of methane emissions from ruminant livestock using a sulfur hexafluoride tracer technique. *Environ Sci Technol.* 1994 Feb 1;28(2):359-62.
- Kang K, Cho H, Jeong S, Jeon S, Lee M, Lee S, Baek Y, Oh J, Seo S. Application of a hand-held laser methane detector for measuring enteric methane emissions from cattle in intensive farming. *J Anim Sci.* 2022 Aug 1;100(8):skab211.
- Kaur U, Malacco VMR, Bai H, Price TP, Datta A, Xin L, Sen S, Nawrocki RA, Chiu G, Sundaram S, Min BC, Daniels KM, White RR, Donkin SS, Brito LF, Voyles RM. Invited review: Integration of technologies and systems for precision animal agriculture—a case study on precision dairy farming. *J Anim Sci.* 2023 Jan 3;101:skad206.
- Kebreab E, Bannink A, Pressman EM, Walker N, Karagianis A, van Gastelen S, Dijkstra J. A meta-analysis of effects of 3-nitrooxypropanol on methane production, yield, and intensity in dairy cattle. *J Dairy Sci.* 2023 Feb;106(2):927-36.
- Korelidou V, Simitzis P, Massouras T, Gelasakis AI. Infrared thermography as a diagnostic tool for the assessment of mastitis in dairy ruminants. *Animals (Basel).* 2024 Sep 16;14(18):2691.
- Leitner S, Ring D, Wanyama GN, Korir D, Pelster DE, Goopy JP, Butterbach-Bahl K, Merbold L. Effect of feeding practices and manure quality on CH₄ and N₂O emissions from uncovered cattle manure heaps in Kenya. *Waste Manag.* 2021 May 1;126:209-20.
- Lewis WH, Lind AE, Sendra KM, Onsbring H, Williams TA, Esteban GF, Hirt RP, Ettema TJG, Embley TM. Convergent evolution of hydrogenosomes from mitochondria by gene transfer and loss. *Mol Biol Evol.* 2020 Feb 1;37(2):524-39.
- Lovendahl P, Difford GF, Li B, Chagunda MGG, Huhtanen P, Lidauer MH, Lassen J, Lund P. Review: Selecting for improved feed efficiency and reduced methane emissions in dairy cattle. *Animal.* 2018 Dec;12(s2):s336-49.
- Mayo LM, Silvia WJ, Ray DL, Jones BW, Stone AE, Tsai IC, Clark JD, Bewley JM, Heersche G Jr. Automated estrous detection using multiple commercial precision dairy monitoring technologies in synchronized dairy cows. *J Dairy Sci.* 2019 Mar;102(3):2645-56.
- McGinn SM, Coulombe JF, Beauchemin KA. Technical note: validation of the GreenFeed system for measuring enteric gas emissions from cattle. *J Anim Sci.* 2021 Mar 1;99(3):skab046.
- McNicol LC, Bowen JM, Ferguson HJ, Bell J, Dewhurst RJ, Duthie CA. Adoption of precision livestock farming technologies has the potential to mitigate greenhouse gas emissions from beef production. *Front Sustain Food Syst.* 2024 Jun 27;8:1414858.
- Minutti JG, Negreya F, Cabrera M, Echeveste JA, Montini J, Gere JI, Cajarville C, Repetto JL, Santana A. Supplementation with a blend of essential oils, tannins, and bioflavonoids in dairy cows. II: Implications for methane emission reduction. *J Dairy Sci.* 2025;108(Suppl_1):423.
- Mizrahi I, Wallace RJ, Morais S. The rumen microbiome: Balancing food security and environmental impacts. *Nat Rev Microbiol.* 2021 Sep;19(9):553-66.
- Montes F, Meinen R, Dell C, Rotz A, Hristov AN, Oh J, Waghorn G, Gerber PJ, Henderson B, Makkar HP, Dijkstra J. Special topics – Mitigation of methane and nitrous oxide emissions from animal operations: II. A review of manure management mitigation options. *J Anim Sci.* 2013 Nov;91(11):5070-94.
- Morgavi DP, Forano E, Martin C, Newbold CJ. Microbial ecosystem and methanogenesis in ruminants. *Animal.* 2010 Jul;4(7):1024-36.
- Munoz C, Letelier PA, Ungerfeld EM, Morales JM, Hube S, Perez-Prieto LA. Effects of pregrazing herbage mass in late spring on enteric methane emissions, dry matter intake, and milk production of dairy cows. *J Dairy Sci.* 2016 Oct;99(10):7945-55.
- Myhre G, Shindell D, Breon FM, Collins W, Fuglestedt J, Huang J, Koch D, Lamarque JF, Lee D, Mendoza B, Nakajima T, Robock A, Stephens G, Takemura T, Zhang H. Anthropogenic and natural radiative forcing. In: *Climate change 2013: The physical science basis. Contribution of Working Group I to the fifth assessment report of the Intergovernmental Panel on Climate Change [Internet].* Cambridge (UK): Cambridge University Press; 2013. p. 659-740 [cited 2026 Jan 20]. Available from: https://www.ipcc.ch/site/assets/uploads/2018/02/WG1AR5_Chapter08_FINAL.pdf.

<https://doi.org/10.17221/116/2025-CJAS>

- NASEM – National Academies of Sciences, Engineering, and Medicine. Nutrient requirements of dairy cattle: Eighth revised edition. Washington, DC: The National Academies Press; 2021. 502 p.
- Norton T, Chen C, Larsen MLV, Berckmans D. Review: Precision livestock farming: Building “digital representations” to bring the animals closer to the farmer. *Animal*. 2019 Dec;13(12):3009-17.
- Olijhoek DW, Lovendahl P, Lassen J, Hellwing ALE, Hoglund JK, Weisbjerg MR, Noel SJ, McLean F, Hojberg O, Lund P. Methane production, rumen fermentation, and diet digestibility of Holstein and Jersey dairy cows being divergent in residual feed intake and fed at 2 forage-to-concentrate ratios. *J Dairy Sci*. 2018 Nov;101(11):9926-40.
- Owen JJ, Silver WL. Greenhouse gas emissions from dairy manure management: A review of field-based studies. *Glob Chang Biol*. 2015 Feb;21(2):550-65.
- Papakonstantinou GI, Voulgarakis N, Terzidou G, Fotos L, Giamouri E, Papatsiros VG. Precision livestock farming technology: applications and challenges of animal welfare and climate change. *Agriculture (Basel)*. 2024 Apr;14(4):620.
- Parker D, Ham J, Woodbury B, Cai L, Spiels M, Rhoades M, Trabue S, Casey K, Todd R, Cole A. Standardization of flux chamber and wind tunnel flux measurements for quantifying volatile organic compound and ammonia emissions from area sources at animal feeding operations. *Atmos Environ*. 2013 Feb;(66):72-83.
- Pereira JE, Zhang J, Jessup R, Jones CS. Editorial: Forage crop improvement for improved livestock production and nutrition. *Front Plant Sci*. 2022 Dec 5;13:1082873.
- Petersen SO, Blanchard M, Chadwick D, Del Prado A, Edouard N, Mosquera J, Sommer SG. Manure management for greenhouse gas mitigation. *Animal*. 2013 Jun;7(Suppl_2):266-82.
- Pfrombeck J, Gandorfer M, Zeiler E, Ettema J. An economic evaluation of sensor-assisted health monitoring in dairy farming using the example of a rumen bolus. *J Dairy Sci*. 2025 Mar;108(3):2573-94.
- Pulina G, Francesconi AHD, Stefanon B, Sevi A, Calamari L, Lacetera N, Dell’Orto V, Pilla F, Marsan PA, Mele M, Rossi F, Bertoni G, Crovetto GM, Ronchi B. Sustainable ruminant production to help feed the planet. *Ital J Anim Sci*. 2017;16(1):140-71.
- Pupo MR, Ferraretto LF, Nicholson CF. Effects of feeding 3-nitrooxypropanol for methane emissions reduction on income over feed costs in the United States. *J Dairy Sci*. 2025 May;108(5):5061-75.
- Roessler R, Chefor F, Schlecht E. Using a portable laser methane detector in goats to assess diurnal, diet- and position-dependent variations in enteric methane emissions. *Comput Electron Agric*. 2018 Jul;(150):110-7.
- Rossi CAS, Grossi S, Dell’Anno M, Compiani R, Rossi L. Effect of a blend of essential oils, bioflavonoids and tannins on in vitro methane production and in vivo production efficiency in dairy cows. *Animals (Basel)*. 2022 Mar 14;12(6):728.
- Rotz CA. Modeling greenhouse gas emissions from dairy farms. *J Dairy Sci*. 2018 Jul;101(7):6675-90.
- Sherman JF, Young EO, Jokela WE, Cavadini J. Impacts of low-disturbance dairy manure incorporation on ammonia and greenhouse gas fluxes in a corn silage-winter rye cover crop system. *J Environ Qual*. 2021 Jul;50(4):836-46.
- Sommer SG, Clough TJ, Balaine N, Hafner SD, Cameron KC. Transformation of organic matter and the emissions of methane and ammonia during storage of liquid manure as affected by acidification. *J Environ Qual*. 2017 May;46(3):514-21.
- Spoliansky R, Edan Y, Parmet Y, Halachmi I. Development of automatic body condition scoring using a low-cost 3-dimensional Kinect camera. *J Dairy Sci*. 2016 Sep;99(9):7714-25.
- Stevenson JS, Hill SL, Nebel RL, DeJarnette JM. Ovulation timing and conception risk after automated activity monitoring in lactating dairy cows. *J Dairy Sci*. 2014 Jul;97(7):4296-308.
- Strapak P, Michakova M, Strapakova E, Neirurerova P, Bujko J. Influence of oestrus on changes of locomotion activity and rumination time in cattle dams. *Acta Fytotech Zootech*. 2021;(24):127-30.
- Studer E, Alsaood M, Steiner A, Becker J. Application note for the use of a wireless device measuring reticular pH under practice conditions in a Swiss dairy herd. *Smart Agric Technol*. 2023 Aug;(4):100170.
- Zsumacher-Strabel M, Potkanski A, Kowalczyk J, Cieslak A, Czauderna M, Gubala A, Jedroszkowiak P. The influence of supplemental fat on rumen volatile fatty acid profile, ammonia and pH levels in sheep fed a standard diet. *J Anim Feed Sci*. 2002;11(4): 577-87.
- Tedeschi LO, Abdalla AL, Alvarez C, Anuga SW, Arango J, Beauchemin KA, Becquet P, Berndt A, Burns R, De Camillis C, Chara J, Echazarreta JM, Hassouna M, Kenny D, Mathot M, Mauricio RM, McClelland SC, Niu M, Onyango AA, Parajuli R, Pereira LGR, Del Prado A, Paz Tieri M, Uwizeye A, Kebreab E. Quantification of methane emitted by ruminants: A review of methods. *J Anim Sci*. 2022 Jul 1;100(7):skac197.
- Thompson LR, Rowntree JE. Invited review: Methane sources, quantification, and mitigation in grazing beef systems. *Appl Anim Sci*. 2020;36(5):556-73.
- Tuytens FAM, Molento CFM, Benaissa S. Twelve threats of precision livestock farming (PLF) for animal welfare. *Front Vet Sci*. 2022 May 27;9:889623.

<https://doi.org/10.17221/116/2025-CJAS>

- USDA. Recommended protocol for using the GreenFeed system to measure gas exchange from cattle as part of the GHG quantification project [Internet]. [place unknown]: United States Department of Agriculture; 2024 Nov 25 [cited 2025 Jan 15]. Available from: <https://www.ars.usda.gov/ARSUserFiles/np212/Greenhouse/GHG%20Research%20Network%20-%20enteric%20GreenFeed%20protocol%20final.pdf>.
- Vakulya G, Hajnal E, Udvardy P, Simon G. In-depth development of a versatile rumen bolus sensor for dairy cattle. *Sensors (Basel)*. 2024 Oct 30;24(21):6976.
- van Breukelen AE, de Haas Y, Aldridge MN, Meijer N, Veerkamp RF. Genetic relationships among methane emissions from breath, dry matter intake, body weight, and milk production traits of Dutch dairy cows. *J Dairy Sci*. 2025 Jul;108(7):7278-96.
- van Lingen HJ, Crompton LA, Hendriks WH, Reynolds CK, Dijkstra J. Meta-analysis of relationships between enteric methane yield and milk fatty acid profile in dairy cattle. *J Dairy Sci*. 2014 Nov;97(11):7115-32.
- Vechi NT, Falk JM, Fredenslund AM, Edjabou ME, Scheutz C. Methane emission rates averaged over a year from ten farm-scale manure storage tanks. *Sci Total Environ*. 2023 Dec 15;904:166610.
- Vladimirov F, Bazaev SO, Khakimov AR. [Comparison of in-rumen systems for monitoring the physiological state of cattle]. *Agric Mach Technol*. 2023 Jun;17(2):35-9. Russian.
- von Soosten D, Meyer U, Flachowsky G, Danicke S. Dairy cow health and greenhouse gas emission intensity. *Dairy*. 2020 Mar 23;1(1):20-9.
- Voulgarakis N, Gougoulis D, Psalla D, Papakonstantinou G, Katsoulos PD, Katsoulis K, Angelidou-Tsifida M, Athanasiou L, Papatsiros V, Christodoulopoulos G. Can computerized rumen mucosal colorimetry serve as an effective field test for managing subacute ruminal acidosis in feedlot cattle? *Vet Res Commun*. 2024 Feb;48(1):475-84.
- Wall E, Simm G, Moran D. Developing breeding schemes to assist mitigation of greenhouse gas emissions. *Animal*. 2010 Mar;4(3):366-76.
- Waters SM, Roskam E, Smith PE, Kenny DA, Popova M, Eugene M, Morgavi DP. International Symposium on Ruminant Physiology: The role of rumen microbiome in the development of methane mitigation strategies for ruminant livestock. *J Dairy Sci*. 2025 Jul;108(7):7591-606.
- Wattiaux MA, Uddin ME, Letelier P, Jackson RD, Larson RA. Invited review: Emission and mitigation of greenhouse gases from dairy farms: The cow, the manure, and the field. *Appl Anim Sci*. 2019 Apr;35(2):238-54.
- Weir C, Duncan ZM, Spore TJ, Corrigan ME, Trigeneyer EC, Ellis WC, Hollenbeck WR, Blasi DA. Evaluation of indwelling pH monitoring technologies. *J Anim Sci*. 2024 Sep 13;102(Suppl_3):350-1.
- Wims CM, Deighton MH, Lewis E, O'Loughlin B, Delaby L, Boland TM, O'Donovan M. Effect of pregrazing herbage mass on methane production, dry matter intake, and milk production of grazing dairy cows during the mid-season period. *J Dairy Sci*. 2010 Oct;93(10):4976-85.
- Xuan T, Zheng T, Li T, Wu B, Li T, Bao W, Qin W. The effects of different doses of 3-NOP on ruminal fermentation parameters, methane production, and the microbiota of lambs in vitro. *Fermentation*. 2024 Aug 23;10(9):440.
- Zaninelli M, Redaelli V, Luzi F, Bronzo V, Mitchell M, Dell'Orto V, Bontempo V, Cattaneo D, Savoini G. First evaluation of infrared thermography as a tool for the monitoring of udder health status in farms of dairy cows. *Sensors (Basel)*. 2018 Mar 14;18(3):862.
- Zimmerman PR, inventor; Zimmerman PR, assignee. System for measuring metabolic gas emissions from animals. United States patent US 5,265,618. 1993 Nov 23.
- Zimmerman PR, Zimmerman RS, inventors; C-Lock Inc, assignee. Method and system for monitoring and reducing ruminant methane production. United States patent US 8,307,785 B2. 2012 Nov 13.

Received: July 28, 2025

Accepted: January 6, 2026

Published online: January 28, 2026