https://doi.org/10.17221/36/2024-CJAS

The impact of parity, litter size and birth weight variations within a litter on piglet pre-weaning performance

Nqobile Lungile Buthelezi[1,2]*, *Bohani Mtileni*[1], *Khathutshelo Agree Nephawe*[1], *Peter Ayodeji Idowu*[1], *Mamokoma Catherine Modiba*[1], *Hezekiel Mpedi*[2], *Takalani Judas Mpofu*[1]

[1]*Faculty of Science, Tshwane University of Technology, Department of Animal Sciences, Pretoria, South Africa*

[2]*Topigs Norsvin-Animal Genetic Centre, Farm Bossemanskraal 538 JR, Bronkhorstspruit, South Africa*

**Corresponding author: nqoshenge11@gmail.com*

Citation: Buthelezi N.L., Mtileni B., Nephawe K.A., Idowu P.A., Modiba M.C., Mpedi H., Mpofu T.J. (2024): The impact of parity, litter size and birth weight variations within a litter on piglets' pre-weaning performance. Czech J. Anim. Sci., 69: 255–268.

Abstract: Sows are selected for their prolificacy. Therefore, assessing difficulties that can be associated with large litters is crucial. This review aims to highlight the factors that can affect reproductive performance such as breed, parity, litter size, environment, and nutrition. Understanding these factors can help producers make informed decisions about sow management and other aspects. Higher within-litter variation is the result of a high number of piglets born. It has been determined that large litter sizes pose a risk to animal welfare in the pig industry. Moreover, large litters bring more piglets with low birth weight leading to high pre-weaning losses. Lighter littermates struggle to get enough colostrum due to the competition with heavy littermates, which lowers their chances of survival causing high mortality. Larger litters may cause greater stress and discomfort of the sow. Therefore, applying stress-free environment and treating the sow with anti-inflammatory compounds may enhance the sow performance and consequent survival of piglets. The litter size is positively correlated with the variation in birth weight within a litter, while the average weight at birth is negatively related to litter size. Producers should therefore focus on potential management strategies that can improve piglet performance, survival and welfare such as proper supervision and adequate colostrum intake. This review highlights the historical development of the pig industry and literature on the variation in piglet weight at birth within a litter, factors influencing weight at birth, and the potential effects on the piglet performance before weaning. Also, the difficulties amongst large litters and possible strategies for managing survival in large litters were also reviewed.

Keywords: colostrum; cross-fostering; littermate; mortality; swine production

The South African pig production sector is regarded as an essential aspect of the country's farming economy (VandeWaal and Deen 2018). This industry consists of both commercial and small-scale pig producers, and it primarily focuses on pork production for local consumption (VandeWaal and

Supported by the Topigs Norsvin SA (No.?) and by the Tshwane University of Technology Postgraduate Scholarship (No?).

© The authors. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0).

https://doi.org/10.17221/36/2024-CJAS

Deen 2018). There are approximately 2 000 commercial pig producers in South Africa across the country. These producers account for most of the country's pork production, with smaller-scale farmers making up the remainder. The South African pig industry is facing some challenges, such as high feed costs, disease epidemics, and opposition to introduced pork products. However, the industry has also made significant strides in recent years, with efforts to improve breeding, genetics, and production techniques (Ahmar et al. 2020). In addition to pork production, pig production is also involved in the production of other pig-derived products, such as bacon, ham, and sausages (Matabane et al. 2018). These products are sold both domestically and internationally, with exports primarily going to other countries in Africa. Overall, the South African pig industry plays a significant part in meeting the protein needs of the population. One of the factors that are responsible for the success and challenges of pig production in South Africa is the pre-weaning performance of piglets.

Piglet pre-weaning performance is an essential factor in the pig industry, as it affects the profitability of pig farming (Grobler et al. 2016). To optimise the maturation and improvement of piglets, it is essential to understand the factors that influence their pre-weaning performance. Previous findings have proven that parity, litter size, birth weight (BW) and variation of piglet weight at birth within a litter significantly influence the piglet performance before weaning (Riddersholm et al. 2021). For instance, sow parity may affect the composition and quantity of milk produced, which may influence the development and survival of her piglets. Similarly to higher litter numbers, smaller litter sizes could not stimulate the sow's milk production adequately, resulting in increased resource competition and slower growth rates. Another crucial element is weight at birth, as piglets with lower BWs are more prone to have health issues and grow at a slower rate (Howe et al. 2020). Pre-weaning performance can also be impacted by variations in BW within a litter since these variations may reflect varying food intake during pregnancy, which could be responsible for variations in growth rates within the litter (Lopez-Soria et al. 2019). Understanding the effects of these factors on pre-weaning performance can help producers make informed decisions about sow management, litter size, and other aspects of pig farming. This can ultimately lead to better outcomes for piglets, improved productivity, and increased profitability for farmers (Blavi et al. 2021).

Therefore, this paper critically reviews the historical improvement in the pig industry, the literature on the variation in BW within a litter, factors influencing it, and the potential effects on piglet performance prior to weaning. Also, the difficulties amongst large litters and possible strategies for managing survival in large litters were also reviewed.

MATERIAL AND METHODS

An electronic search in Google Scholar, PubMed, Science Direct and SciELO databases was used to gather information for the review. The keywords that were used include colostrum, littermate, mortality, cross-fostering, and swine production. The words were used to search for as many articles as possible. A total of 109 articles were selected. During the research, only articles discussing the pre-weaning performance of piglets according to parity, litter size and BW variations were selected.

Birth weight: A major factor for weaning weight

Piglet weight at birth is the most significant factor of weaning weight, the main component of weaning capability. Even though it is challenging to manipulate, there are several strategies to improve it (Yuan et al. 2015). The most common way to improve piglet viability is by increasing the sow nutrition settings in the last 21 to 28 days of maternity, which is usually 2.6 kg in primiparous sows and 2.8 kg in multiparous sows. Although the effect on weight at birth is usually minor, this method was found to have a positive effect on piglet viability. Recent research indicates that adding oil to the diet (about 5% of oil) may result in fewer stillborn piglets, better endurance for smaller piglets, and improved weaning weights (Swiątkiewicz et al. 2020; Vodolazska and Lauridsen 2020). To enhance BW, accurate feed levels and attention to sow health during the breeding cycle are essential (Vodolazska and Lauridsen 2020).

In practice, weight at birth is the common sign of piglet's ability to live up to weaning (Feldpausch et al. 2016; Tucker et al. 2021). Several authors

https://doi.org/10.17221/36/2024-CJAS

(Magnabosco et al. 2016; Zotti et al. 2017; Romero et al. 2022) have looked at the long-term effect of light weight at birth on piglet endurance and maturation, but their results on the piglet capacity to make up for lost time and catch up with their littermates in terms of growth have been inconsistent. Although some have claimed that low BW piglets are more likely to exhibit poor pre-weaning maturation and lighter body weight at weaning, it has been shown that piglets with light weight at birth can counterbalance in post-natal development (Howe et al. 2020). In addition, piglets with lower BW have slower lifespan growth rates and more life before market weight than piglets with higher BW (Howe et al. 2020).

According to Nuntapaitoon et al. (2018), other than birth weight as the indicator of pre-weaning growth and survival, intra-uterine growth restriction (IUGR) can also affect the performance of piglets. While some small foetuses are naturally small but healthy, some are the outcome of placental inability, which can lead to IUGR piglets. These neonates contribute significantly to pre-natal death in numerous species. Delayed foetal growth can be categorized into two types: unbalanced and balanced. Unbalanced growth refers to a body that is extremely smaller than the head, while balanced growth affects the head in part to the rest of the body. As per Amdi et al. (2017) and Huting et al. (2018), the most common type of IUGR piglet is asymmetrical, which is characterized by a dolphin-like head shape.

The IUGR pigs develop in a dilatory way compared to their littermates, with a reduced vigour reserve, immature organs, and anosmic gastrointestinal system development (Hales et al. 2014).

Factors influencing the birth weight and within-litter variation

Several factors affect weight at birth and within-litter variation in pig production, including breeds, parity, temperature, litter size, and management, among others. These variables all play a role in within-litter weight differences. Therefore, it is crucial to concentrate on pre-service and gestation periods to enhance oocyte and conceptus development uniformity, minimize variation during implantation, and improve newborn piglet weight (Yuan et al. 2015).

Breed/strain. The pig breed is a crucial factor in determining their reproductive performance, with both purebred and crossbred sows being used by producers in their efforts to increase productivity (Knecht et al. 2015). According to Getahun et al. (2019), crossbreeding is particularly beneficial due to the hybrid vigour it imparts, resulting in positive effects on traits like fertility and survivability, which have low heritability. This, in turn, translates to improved conception rates, birth weights, litter sizes, and weaning rates. Moreover, crossbred dams also exhibit maternal heterosis, which enhances piglet weight, including birth weight, within-litter variation, and weaning weight, as well as litter size (Pedersen et al. 2019).

According to research, breed is a significant factor that affects reproductive performance in pigs. To address this issue, crossbreeding has been identified as an effective method to improve generative production because of the low heritability of reproductive traits (Hagan and Etim 2019). Furthermore, a larger number of piglets born is an indication of the inherent perspective of the breed's generative biology, as well as the cros bred's ability to withstand environmental factors like weather conditions, nutritional factors, and other similar challenges (Hagan and Etim 2019). The most popular breeds used in crossbreeding are Large White (LW) and Landrace (L) breeds due to their great performance in the pig industry. LW are normally selected as dam lines in breeding programs because of their outstanding number of piglets born and healthier litter with high survivability (Rothschild and Ruvinsky 2011), while L pigs are commonly used as a sire line because of their high fertility (Pedersen et al. 2019). The LW and L produce a high number of born alive piglets and increased litter weight at birth (Amatucci et al. 2022). Nevertheless, the larger number of piglets born is associated with light weight piglets at birth and the increase in within-litter variation, consequently, increases pre-weaning mortality. The use of crossbreds improves conception rates leading to a higher pregnancy rate, farrowing and higher litter size (Hagan and Etim 2019) although this can result in high pre-weaning mortality. Piglet survival rates are due to the mothering ability of the sow (Edwards and Baxter 2015), thus high litter size at birth may lead to negative effects when the dam receives insufficient diet to make adequate milk in the early period after parturition.

https://doi.org/10.17221/36/2024-CJAS

Parity. The impact of parity on piglet weight at birth is the cause of the large number of piglets born, which results in greater variability (Zindove et al. 2013). Litter size tends to increase up to parity 3 and then it declines with the dam's age. The increase in age leads to the increase of stillborn piglets, and birth weight becomes more variable. Older sows with higher parity tend to have fewer uniform litters and a higher proportion of light weight piglets at birth (Yuan et al. 2015; Charneca et al. 2023). In contrast, piglets born to primiparous sows tend to have much uniform birth weights (Yuan et al. 2015). Litter size has a positive relationship with parity, with the highest levels typically reached between parity 3 and 5 (Freyer 2018). However, the negative impact of parity on litter uniformity is possibly due to its association with birth weight variation within a litter, which tends to rise with larger litter sizes (Quesnel et al. 2014). Moreover, the birth weight variation tends to be higher in older parities after accounting for the total number of pigs born (Wientjes et al. 2013). Some researchers have suggested that the change in litter similarity in old sows may be due to the decline in oocyte quality that accompanies aging (Yuan et al. 2015), indicating that parity can have an important impact on litter uniformity in terms of BW.

Temperature. Pigs have a predisposition to heat stress as they do not sweat very much and rely on evaporative cooling according to Boddicker et al. (2014). Unfortunately, modern genetic selection has exacerbated this issue, since pigs are selectively bred for excessive lean tissue growth, increasing their basal heat production, as noted by Johnson et al. (2015). When pregnant sows face heat stress, it can lead to changes in the growth, body composition, and metabolic function of piglets, with the severity of these changes dependent on the duration and intensity of stress, as outlined by Boddicker et al. (2014) and Johnson et al. (2015). These changes are brought about by alterations in metabolism and uterine blood flow, leading to differences in skeletal and intestinal growth, as well as hormonal and metabolic profiles. This can have a negative influence on piglet growth and performance, as highlighted by Dhakal et al. (2013) and Boddicker et al. (2014). Additionally, Rutherford et al. (2013) suggested that uteroplacental vascularity differences may affect some piglets more than others, leading to variations in weight and growth.

Litter size. The breeding of highly productive sows causes an increase in the number of pigs born (Koketsu et al. 2017). However, a large number of piglets born causes a decrease in weight at birth and an increased variation within a litter (Charneca et al. 2021; Peltoniemi et al. 2021; Nam and Sukon 2022). Low BW piglets are disposed to perinatal death (Nam and Sukon 2020) and have a lower endurance rate and lower growth rate (Riddersholm et al. 2021; Nam and Sukon 2022). Increased litter size leads to a huge proportion of light piglets having insufficient body strength, resulting in sensitivity to cold (Kitkha et al. 2017). Besides, they struggle to compete with their littermates for adequate colostrum and milk consumption, leading to higher chances of death in the early days after birth (Kitkha et al. 2017).

Maintaining an optimal litter size is crucial in the pig industry, and as such, reducing litter size is not encouraged due to its impact on profits (Zindove et al. 2013; Marandu et al. 2015). A larger litter requires a plenty of nutrients before and after parturition, which can be challenging to provide (Baxter et al. 2013). However, improved litter size can lead to increased milk production, which then requires more energy from the sow to sustain production due to suckling intensity (Muns et al. 2023). If the sow experiences a body condition loss, this can result in compromised foetal development, leading to higher variation within a litter (Rutherford et al. 2013; Wientjies et al. 2013). Furthermore, the sow's ability to balance increased feed consumption is reduced, and low consumption can affect generative hormone levels, specifically those with an impact on follicle development (Langendijk 2021). This can significantly impact on post-lactation production by delaying follicular development and oestrus (Bierhals et al. 2012; Soede et al. 2015). To alleviate the stress of large litters, re-homing piglets is practised instead of selecting against it (Pedersen et al. 2019).

Challenges associated with larger litter. In the pig farming industry, sows are carefully chosen and bred for their ability to produce larger litters, ultimately resulting in a higher number of pigs being sold each year (Yuan et al. 2015). However, as producers strive to increase litter size, it is essential to consider potential issues that may arise and determine if these challenges could be successfully controlled or need medical interference (Ward et al. 2020). As litter size increases, sows

https://doi.org/10.17221/36/2024-CJAS

face various obstacles, including higher rates of preweaning and post-weaning mortality, and developmental concerns (Tucker et al. 2021).

Intrauterine crowding and its effect on piglet improvement. Even though the sows can give birth to larger litters, it is crucial to keep in mind that the uterine space and body fluid provision are restricted assets, as supported by research conducted by Pardo et al. (2013), Madsen and Bee (2015), and Langendijk et al. (2016). Typically, a sow will begin pregnancy with approximately 15–20 viable embryos, with only 9–13 eventually developing into live piglets, as Greenwood et al. (2014) have shown. Despite this, it is worth noting that commercial production has seen litters greater than 16 piglets become increasingly common, as observed by Fijn et al. (2016) and Roelofs et al. (2019). In cases where the litter size is larger, the sow uterus can become crowded with embryos, leading to embryonic competition. This competition can occur as the first implanted embryos limit the process of joining embryos, and the contest rises with each thriving immature attachment, as indicated by Ward et al. (2020).

Moreover, when the uterus surpasses its normal capacity, each additional littermate results in a decline in individual foetal growth (Ward et al. 2020). As a result, a larger litter size is strongly connected with a substantial balance of piglets being born with low weight (less than 1.0 kg), as noted by Madsen and Bee (2015) and Schmitt et al. (2019). These piglets with light weight at birth have a higher surface area to volume ratio, they are vulnerable to poor viability, weakened immunity, hypothermia, and hypoglycaemia in the early hours of birth (Ward et al. 2020). Thus, light-weight piglets have a higher hazard of mortality before weaning compared to their average-weight counterparts (Zeng et al. 2019).

Additionally, intrauterine crowding can impede the physiological development of the foetus during gestation (Amdi et al. 2013). The piglets affected by IUGR not only exhibit a disproportionate physical appearance at birth, including the head shape reminiscent of a dolphin, as noted by Amdi et al. (2013), Hales et al. (2014), and Chand et al. (2022), but they also suffer from immature intestinal development, leading to metabolic compromise (Engelsmann et al. 2019). Inflammatory and metabolic profiles are further disturbed in these piglets, as highlighted by Huang et al. (2019) and Wen et al. (2022). Piglets exposed to intrauterine restriction exhibit insufficient milk consumption and hepatic glycogen, resulting in higher morbidity and mortality rates, as well as hypothermia (Magnabosco et al. 2015; Vallet et al. 2016).

Higher incidence of piglets being crushed by a sow. Researchers have reached a consensus that the primary reason for early piglet death is the overlaying by the sow. Contributing factors include pig breed and litter size, increased sow parity, bad mothering ability, and reduced birth weight. Additionally, sows or gilts experiencing stress and irritation during the periparturient period are restless and maximize the possibility of crushing, especially if the sow immediately lies down. According to Ward et al. (2020), crushing can potentially be a scheme to minimize maternal investment in larger litters, since the prevalence of sows crushing any piglets was greater for prolific sows. Weak piglets and those with lower viability are more susceptible to being crushed (Muns et al. 2017).

The welfare of piglets plays a crucial role in their survival until they are weaned. Research shows that pre-weaning survival is positively associated with birth weight uniformity within a litter from a genetic perspective (Charneca et al. 2023). Piglet survival is linearly correlated with the coefficient of variation for birth weight within a litter, while the relation between mean birth weight and mortality is curvilinear (Pardo et al. 2013; Yuan et al. 2015). Uniformity in litter birth weight is also linked to higher survival rates and improved welfare (Zindove et al. 2013; Charneca et al. 2021). According to Marandu et al. (2015), individual weight at birth has a more significant impact on piglet survival compared to relative birth weight. Additionally, male piglets have a greater mortality rate caused by higher basal cortisol levels, making them more susceptible to stress. Higher within-litter variability increases the likelihood of mortality before weaning, as smaller piglets must contend with heavier ones for resources. This contest leads to lighter piglets remaining closer to the dam for a longer time, increasing their risk of getting crushed (Yuan et al. 2015). It has been determined that litters with a reasonable size of 10 to 14 piglets are more likely to survive up to weaning (Marandu et al. 2015).

Potential strategies for managing survival in large litter

The quantity and quality of piglets that are weaned and their subsequent performance in a later phase

https://doi.org/10.17221/36/2024-CJAS

of production are affected by the level of attention and management given in the farrowing housing. The best management strategies to reduce piglet mortality are to increase birth weight and improve heating (Harper and Bunter 2024), giving the piglet more energy to stay far from the sow (Ward et al. 2020). Birth weight can be built through exclusive breeding, nutrition (Renteria Flores et al. 2021), and early fostering (Zhang et al. 2021). All these management strategies include managing integrated piglets and sow management, colostrum intake, environment, nutrition, and herd parity structure management.

Integrated piglets and sow management. Piglets that lack uniformity in size require more intensive management, which can lead to increased costs including feeding (Yuan et al. 2015; Tian et al. 2016). For instance, smaller piglets may need more time to get the market weight compared to larger littermates, which can impact on the overall cost of production. To address this, piglet transfer is a popular practice in pig production to balance litter numbers and minimize the influence of BW variation (Heim et al. 2012). However, this process needs extra effort and time. To ensure the survival and proper colostrum sharing of small piglets, cross-fostering should take place as soon as possible following the birth, preferably within 24 h or sooner after farrowing. In cases where this is not feasible, split suckling can facilitate colostrum sharing in larger litters (Romero et al. 2023). Mortality rates can increase if piglet numbers are higher than the sow rearing ability, making fostering a necessary practice across the industry. This helps to re-home piglets associated with increased sow prolificacy in relation to the number of available useful teats (Vande Pol et al. 2021; Oliviero 2023). Additionally, fostering can guarantee litter uniformity by reducing variability in weaning weight, which can have subsequent impacts on slaughter management (Van den Bosch et al. 2019). However, the process of relocating a piglet from one birth mother to another can be challenging. It can result in changes to the surroundings, littermates, and sow, which can be disadvantageous for the moved piglet and may irritate the litter to which it is transferred (Baxter et al. 2013; Schmitt et al. 2019).

In the typical all-in-all-out system, variation poses a challenge, as pigs must be sorted by weight rather than by age. However, this system has the benefit of preventing disease transmission by halting the infection and inhibiting the accumulation of pathogens through facility disinfection between groups. Additionally, it enhances feed productivity and everyday strength, ultimately resulting in fewer days to merchandising and decreased feed prices (Aymerich et al. 2020). The increased uniformity achieved through this system also allows for easier management and lower mortality rates, and consistent size and weight of the final product, and also quality attributes such as colour, marbling, and drip loss, prove advantageous for both retailers and consumers alike (Roche et al. 2010). It is worth noting that variation in pig weight and shape can result in controlling troubles and product regularity matters which would be detrimental (Zindove et al. 2013; Mulder and Rashidi 2015). Given that profitability is tied to slaughter and management procedures, it is prudent to opt for sows that give more homogeneous litters (Zindove et al. 2013).

Colostrum consumption. Newborn piglets have lower energy levels and fewer protective antibodies, making them reliant on colostrum from the sow immediately after birth to provide them with energy and protection against harmful pathogens like viruses and bacteria (Rasmussen et al. 2016). Foremilk, which is the first milk produced by the sow, is an essential source of nutrients, resistance, and energy for the piglet (Silva et al. 2019). Immediately after giving birth, it is secreted from the udder and its composition changes within a few hours to match that of sow milk (Silva et al. 2019).

During birth, piglets experience a sudden shift in their energy supply as they begin to feed. In addition, the environment in which they are born is usually quite cold. Due to their high surface area to volume ratio and wet skin, new-born piglets lose heat quickly, causing a sharp drop in their body temperature (Theil et al. 2014). With < 2% body fat and high vigour demands, piglets get into a negative energy balance soon after birth (Theil et al. 2014). Therefore, it is crucial for piglets to consume colostrum shortly after birth to maintain their energy balance. Colostrum is most beneficial when obtained from a piglet's birth sow. Recently, it has been revealed that colostrum contains certain aspects of immunity (cell-mediated immunity) that are only present in piglets that have been preserved on their original dam for more than 12 hours (Bandrick et al. 2014).

Research has proved that light birth weight piglets (800–1 200 g) have an over 89% chance of surviving

https://doi.org/10.17221/36/2024-CJAS

until weaning if they receive about 200 ml of colostrum (Moreira et al. 2017). Declerck et al. (2016) also reported a similar observation, finding that a correlation exists between inadequate colostrum ingestion and decreased survival before weaning, with the higher effects in piglets with lower weight at birth. Furthermore, colostrum has a significant impact on the development of the neonatal gut (Ward et al. 2020). Colostrum provides essential nutrients for vigour, growth, and development to neonates for their transition from the placental supply of substantial nutrients such as glucose and amino acids after birth (Sangild et al. 2021). Additionally, colostrum offers immune protection that assists the element of neonatal innate immunity up to the maturation of adaptive immunity to the environment (Sangild et al. 2021).

In managing larger litters, it is important to prioritize the colostrum ingestion of lower weight pigs. A viable approach is the split suckling technique, as demonstrated by Alexopoulos et al. (2018) and Tang et al. (2023). This technique includes dividing the piglets into different groups, with the heavier piglets temporarily separated to allow smaller piglets' access to the udder and benefit from colostrum intake, a strategy supported by Kirkden et al. (2013) and Vandaele et al. (2020). The ultimate purpose of split suckling is to minimize competition for colostrum (Morton et al. 2019).

Environmental management. The viability and survival of embryos may be largely dependent on the quality of the oocyte, which is in turn affected by various factors including maternal situation, nourishment, temperature, and sow age (Swinbourne et al. 2014). The number of pigs per litter is given by the number of fertilized and released gametes that evolve into viable offspring. According to Swinbourne et al. (2014), the smaller oocyte size during temperate times can lead to smaller litters due to the heat impact on hormone production and oocyte competence. Heat stress during gestation can be particularly detrimental in the first half of the time (Boddicker et al. 2014). Fibre growth in pigs is limited to days 35 and 90 of maternity, and heat stress during early gestation can negatively impact on the primary fibre growth, leading to suboptimal performance (Boddicker et al. 2014). Johnson et al. (2015) also observed that heat-stressed piglets had smaller organs, especially the liver, which plays a part in energy production for the first few days of life because of its high glycogen concentration (Theil et al. 2014). These factors can lead to variable BWs within litters as they affect individual piglets differently *in utero*.

During two hours after farrowing, immature piglets are particularly vulnerable to the physiological state. The type of flooring in their environment can have an impact on their body temperature, as metal slats can lower their rectal temperature when compared to solid concrete floors (Pedersen et al. 2016). To optimize piglet performance, the plastic-coated expanded metal flooring is a better option than concrete slats (Ramirez et al. 2022). The most effective way to manage piglet environments is to maintain two distinct microclimates – one for piglets kept at a temperature of 30–35 °C, and the other for sows kept between 18 and 20 °C. This approach reduces utility costs and minimizes the sow heat stress compared to the entire farrowing house, while still supporting piglet needs (Ramirez et al. 2022). To create an ideal microclimate for piglets, it is important to consider the type of supplemental heat source and adequate space allocation (Ramirez et al. 2022).

Nutrition. During pregnancy, the developing litter is granted high precedence for vigour and nutrients, though the demand is quite small. As a result, it is improbable that ingestion levels during maternity will have an impact on litter size under normal commercial conditions (Harper and Bunter 2024). Additionally, it has been found that excessively high vitamin levels in herds with normal diets do not increase the litter size. However, experiments conducted in Canada have revealed that severely limited feed consumption in any week of suckling can affect ovulation rate, embryo survival, and ultimately litter size (Theil et al. 2014; Renteria et al. 2021).

According to Liu et al. (2020), feed consumption of sows during gestation can also affect piglet birth weight similarity. The intake of cycling gilts at or below the maintenance level, as opposed to *ad libitum*, has been shown to decrease oocyte maturation (Swinbourne et al. 2014). This difference in gametocyte growth rates inside the uterus, potentially due to uneven nutrient accessibility to the foetuses, can result in variations in foetus and piglet weight. The litter-of-origin consequence may also impact on the future generative productivity of gilts, as larger litters can decrease colostrum availability and hinder neonatal uterine gland development (Vallet et al. 2016).

https://doi.org/10.17221/36/2024-CJAS

To increase ovulation rate and litter size, it is recommended to feed gilts quality diet *ad libitum* at least 10 d prior to breeding. However, feeding gilts a higher ration (more than 2.5 kg feed/day) in 48 h to 72 h after breeding can minimize the embryo survival (Theil et al. 2014; Renteria et al. 2021) and ultimately the litter size.

Herd parity structure management. In order to consistently produce the highest quality piglets, it is essential for the herd parity structure to remain stable. This can only be achieved through a consistent influx of gilts into the herd, maintaining a greater number of females in the 3–6 parity range, and strictly culling based on age after 7–8 parities. Failing to maintain a balanced parity structure due to the uneven gilt section, high removal rates among immature females or poor culling policies can create unwanted variations in mean litter size and birth weight over time. This, in turn, can have several counter implications for the maturation and efficiency of pigs from weaning to market (Marandu et al. 2015). Both piglet survival and litter weaning weight decrease after the sow approaches the third or fourth parity (Zeng et al. 2019). The likely explanation for reduced piglet survival is enhanced sow unwellness stress acting to ease milk yield (Tummaruk and Sang-Gassanee 2013) and poor teat availability (Tummaruk and Sang-Gassanee 2013). Furthermore, older sows are more disposed to lameness (Li and Gonyou 2013) and this can lead to a high risk of crushing (Pluym et al. 2013), causing a high mortality rate.

The productivity of breeding herds is mostly assessed by the number of pigs weaned per sow per year (Koketsu et al. 2017). However, pigs weaned per sow per year are a good indicator of the herd ratio, but not of sow longevity and piglet quantity (Koketsu et al. 2017). As the number of pigs born alive increases, the birth weight decreases, and small piglets fail to consume sufficient colostrum from the sow. Therefore, lighter birth weight and insufficient colostrum consumption result in higher pre-weaning mortality (Declerck et al. 2016). The number of pigs born per sow per year is influenced by the farrowing rate and culling intervals because of unproductive days caused by abortion in commercial herds (Iida and Koketsu 2015; Declerck et al. 2016).

The fertility and prolificacy of sows depend on the herd sow management including gilt development program, time of insemination, farrowing and lactation management and culling strategies (Koketsu et al. 2017). Kaneko et al. (2013) indicated that gilts inseminated immediately after oestrus detection have higher farrowing rates than those with delayed insemination. According to Koketsu and Iida's (2020) findings, the recommended culling interval for mated gilts and sows was exceeded by a minimum of 30 days. As a result, producers should employ echography reproduction of ovaries in gilts to determine the timing of insemination in connection with ovulation, as suggested by Koketsu et al. (2017).

Birth weight within-litter variation management. Managing the low uniformity piglets can be challenging. For example, smaller piglets may require an additional time to get to the market weight when compared to their larger littermates, leading to an increase in feeding costs (Yuan et al. 2015; Tian et al. 2016). To address this issue, piglet relocation is a popular practice in pig farming, which involves merging the litter size to the number of teats and mitigating the impacts of birth weight variation (Vande Pol et al. 2021). However, this process requires extra labour and time. In the model all-in-all-out system, sorting pigs by weight, not by age, is crucial due to variation. This system helps prevent disease transmission as facilities are disinfected between groups.

Consequences of birth weight within-litter variation

The consequences of variations in BW within a litter can result in lower pre-weaning growth performance, as well as impact on the vitality and survival of piglets, as noted by Yuan et al. (2015) in their study. This dispute about growth performance can be attributed to both the disparity in milk intake and the efficiency of nutrient utilization among littermates. Studies have confirmed that piglets with low weight at birth have a better chance of thriving during the pre-weaning stage if raised alongside piglets of the same weight, instead of the heavier ones (Huting et al. 2018; Vande Pol et al. 2021).

Pre-weaning growth performance. Studies have shown that there is a relation between birth weight variations in piglets and their pre-weaning growth performance within litters. These variations can be largely attributed to differences in colostrum consumption and nutrient utilization efficiency

https://doi.org/10.17221/36/2024-CJAS

among littermates (Yuan et al. 2015). The consumption of colostrum, which is important for piglet growth and development, is influenced by piglet vitality at birth and competition among littermates for access to teats. The teat order constituted within the first 24 h after birth tends to persist. Studies have also found that heavier pigs have a high likelihood of endurance compared to their lighter littermates because they can consume larger amounts of colostrum. This is so because the ability to suckle more colostrum is linked to birth weight (Declerck et al. 2016; Tucker et al. 2021). Furthermore, higher BW and colostrum intake are also associated with better growth performance, as evidenced by studies conducted by Declerck et al. (2016), Le Dividivh et al. (2017), and Vodolazska et al. (2023).

Vitality and survival of piglets. Piglet vitality refers to the piglet capacity to survive from birth to weaning, as determined by its survival rate during this period (Tucker et al. 2021). Vigour and endurance traits are affected by additional genetic parameters (such as behaviour, vigour, immunity), maternal genetic parameters (such as behaviour, milk choice, and amount), uterus choice, average litter (such as litter size), and several situation effects (such as temperature, stress, and difficulties during farrowing, as well as assistance with colostrum intake), which are difficult to statistically identify (Heub et al. 2019). Birth weight is the primary determinant of piglet survival, and an alternative trait to consider when breeding for improved piglet endurance due to its high heritability (Heub et al. 2019).

Once body reserves and thermoregulatory ability are ensured, the main factor that influences piglet endurance is the vigour of the piglet instantly after parturition. Neonatal vitality can be measured through behavioural findings, such as the period to get to the udder, and physiological parameters, including the muscle tone. Piglet vigour varies significantly among littermates, and this change is influenced by the degree of hypoxia at birth (Farmer and Edwards 2020). The last-born piglets typically experience hypoxia. Neonatal asphyxia results in critical acidosis and has severe negative impacts on the function of essential organs. Decreased total vigour, weak thermoregulation, poor inactive transfer of immunity, weak performance, and higher susceptibility to infections are crucial secondary problems related to neonatal asphyxia and acidosis (Bailey et al. 2020). Low birth weight piglets have an enhanced risk of death and decreased weight gain during suckling. For instance, piglets with < 1 kg weight at birth are most likely to die prior to weaning (Marandu et al. 2015).

PROSPECTS AND OPPORTUNITIES

The study of the impacts of parity, litter size, birth weight, and within-litter birth weight variation on pre-weaning performance in piglets has significant implications for the future of pig farming. By understanding how these factors impact on piglet growth, health, and survival, producers can implement strategies to improve pre-weaning performance and overall productivity (Schinckel et al. 2010).

One opportunity for future research is the development of breeding programs that select for traits that promote better pre-weaning performance. This could include selecting sows with higher milk production or piglets with better growth rates and disease resistance. With the growing importance of animal welfare in agriculture, there is also a potential for research on how to optimize pre-weaning performance while ensuring the health and well-being of sows and piglets (Serenius et al. 2019).

Another opportunity is the development of innovative management practices that can help mitigate the negative effects of factors such as large litter sizes or low birth weights. For example, the use of foster sows or cross-fostering techniques can help redistribute piglets to ensure that they will receive adequate nutrition and care. Advances in technology and precision farming could also help improve the monitoring and management of sow and piglet health and nutrition (Wellock et al. 2014). In addition, there is a potential for the pig industry to expand and diversify its product offerings beyond pork, such as producing pig-derived pharmaceuticals or other value-added products. This could provide new revenue streams and opportunities for pig farmers.

CONCLUSION

Based on the literature reviewed, piglets born with lower weight face a higher risk of death, especially when they have not received sufficient colostrum intake. Poor pre-weaning performance is often caused by low BW and large litter size,

https://doi.org/10.17221/36/2024-CJAS

which must be managed promptly. While economic considerations drive present management practices, a more ethical approach is necessary. Instead of focusing solely on the number of pigs weaned, producers must also consider the number of piglets that died to reach that figure. The endurance of smaller piglets during weaning is particularly challenging due to the enhanced risk of low BW and weight variation within the litter. Colostrum is vital for their survival, and piglets should receive it within 24 h of birth from their birth sow. When the sow's colostrum production is insufficient, fostering to another farrowing sow may be necessary. Split suckling can provide smaller piglets with a chance to suckle, and efforts should be made to ensure the even colostrum ingestion among litters. Piglet transfer should occur within 12–24 h after birth, and alternative schemes, including the use of foster sows, should be engaged if piglet movement must be avoided.

Acknowledgement

We thank Topigs Norsvin South Africa and the Tshwane University of Technology for their contribution to the study.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

Ahmar S, Gill RA, Jung KH, Faheem A, Qasim MU, Mubeen M, Zhou W. Conventional and molecular techniques from simple breeding to speed breeding in crop plants: Recent advances and future outlook. Int J Mol Sci. 2020 Apr 8;21(7):2590.

Alexopoulos JG, Lines DS, Hallett S, Plush KJ. A review of success factors for piglet fostering in lactation. Animals (Basel). 2018 Mar 9;8(3):38.

Amatucci L, Luise D, Correa F, Bosi P, Trevisi P. Importance of breed, parity, and sow colostrum components on litter performance and health. 2022 May 10;12(10):1230.

Amdi C, Krogh U, Flummer C, Oksbjerg N, Hansen CF, Theil PK. Intrauterine growth restricted piglets defined by their head shape ingest insufficient amounts of colostrum. J Anim Sci. 2013 Dec;91(12):5605-13.

Amdi C, Jensen LL, Oksbjerg N, Hansen CF. Supplementing new-born intrauterine growth restricted piglets with a bolus of porcine colostrum raises rectal temperatures one degree Celsius. J Anim Sci. 2017 Jul;95(7):2968-76.

Aymerich P, Soldevila C, Bonet J, Gasa J, Coma J, Sola-Oriol D. Increasing dietary lysine impacts differently growth performance of growing pigs sorted by body weight. Animals (Basel). 2020 Jun 13;10(6):1032.

Bailey EJ, Frolova AI, Lopez JD, Raghuraman N, Macones A, Cahill AG. Mild neonatal acidemia is associated with neonatal morbidity at term. Am J Perinatol. 2021 Aug; 38(S01):e155-e61.

Bandrick M, Ariza-Nieto C, Baidoo SK, Molitor TW. Colostral antibody-mediated and cell-mediated immunity contributes to innate and antigen-specific immunity in piglets. Dev Comp Immunol. 2014 Mar;43(1):114-20.

Baxter EM, Rutherford KM, D'eath RB, Arnott G, Turner SP, Sandoe P, Moustsen VA, Thorup F, Edwards SA, Lawrence AB. The welfare implications of large litter size in the domestic pig II: Management factors. Animal Welfare. 2013 May 22;(2):219-38.

Bierhals T, Magnabosco D, Ribeiro RR, Perin J, DA Cruz RA, Bernardi ML, Wentz I, Bortolozzo FP. Influence of pig weight classification at cross-fostering on the performance of the primiparous sow and the adopted litter. Liv Sci. 2012 Apr;146(1):115-22.

Blavi L, Sola-Oriol D, Llonch P, Lopez-Verge S, Martín-Orue SM, Perez JF. Management and feeding strategies in early life to increase piglet performance and welfare around weaning: A review. Animals (Basel). 2021 Jan 25; 11(2):302.

Boddicker RL, Seibert JT, Johnson JS, Pearce SC, Selsby JT, Gabler NK, Lucy MC, Safranski TJ, Rhoads RP, Baumgard LH, Ross JW. Gestational heat stress alters postnatal offspring body composition indices and metabolic parameters in pigs. PLoS One. 2014 Nov 10;9(11):e110859.

Chand KK, Pannek K, Colditz PB, Wixey JA. Brain outcomes in runted piglets: A translational model of fetal growth restriction. Dev Neurosci. 2022;44(4-5):194-04.

Charneca R, Freitas A, Nunes J, Le Dividich J. Effects of the mean weight of uniform litters on sows and offspring performance. Animals (Basel). 2023 Oct 4;13(19):3100.

Charneca R, Nunes JT, Freitas A, le Dividich J. Effect of litter birth weight standardization before first suckling on colostrum intake, passive immunization, pre-weaning survival, and growth of the piglets. Animal. 2021 Apr;15(4):100184.

Declerck I, Dewulf J, Sarrazin S, Maes D. Long-term effects of colostrum intake on piglet mortality and performance. J Anim Sci. 2016 Apr;94(4):1633-43.

Dhakal S, Auckland C, Huang Y, Ambros B, Duke T, Wilson DG, Foxcroft G, Harding J. Uterine spaciousness

https://doi.org/10.17221/36/2024-CJAS

during embryo and fatal development in multiparous sows improves birth weight and postnatal growth performance. Liv Sci. 2013;153(1):154-64.

Edwards SA, Baxter EM. Piglet mortality: Causes and prevention. In: Farmer C, editor. The gestating and lactating sow. Wageningen, Wageningen Academic, 2015; p. 253-78.

Engelsmann MN, Hansen CF, Nielsen MN, Kristensen AR, Amdi C. Glucose injections at birth, warmth and placing at a nurse sow improve the growth of IUGR piglets. Animals (Basel). 2019 Aug 2;9(8):519.

Farmer C, Edwards SA. The neonatal pig: Developmental influences on vitality. In: Farmer C, editor. The suckling and weaned piglet. Wageningen, Wageningen Academic, 2020; p. 649-53.

Feldpausch J, Jourquin J, Bergstrom J, Bokenkroger C, Nelssen JL, Ritter M, Davis DL, Gonzalez JM. Birth weight threshold for identifying piglets at-risk for pre-weaning mortality. Transl Anim Sci. 2019 Jun 5;3(2):633-40.

Fijn L, Antonides A, Aalderink D, Nordquist RE, Van der Staay FJ. Does litter size affect emotionality, spatial learning, and memory in piglets? Appl Anim Behav Sci. 2016 Mar;178(1):23-31.

Freyer G. Maximum number of total born piglets in a parity and individual ranges in litter size expressed as specific characteristics of sows. J Anim Sci. Technol. 2018 May 21; 60(1): 13.

Getahun D, Alemneh T, Akeberegn D, Getabalew M, Zewdie D. Importance of hybrid vigor or heterosis for animal breeding. Biochem Biotechnol Res. 2019 Jul;7(1):1-4.

Greenwood EC, Plush KJ, van Wettere WH., Hughes PE. Hierarchy formation in newly mixed, group housed sows and management strategies aimed at reducing its impact. App Anim Behav Sci. 2014 Nov;160(1):1-11.

Grobler J, Van Schalkwyk SJ, Swatson HK. The South African pig industry, a profile. S Afr J Anim Sci. 2016;46(2):123-34.

Hagan JK, Etim NN. The effects of breed, season, and parity on the reproductive performance of pigs reared under hot and humid environments. Trop Anim Health Prod. 2019 Feb;51(2):411-8.

Hales J, Moustsen VA, Nielsen MBF, Hansen CF. Higher pre-weaning mortality in free farrowing pens compared with farrowing crates in three commercial pig farms. Animal. 2014 Jan;8(1):113-20.

Harper J, Bunter KL. Improving pig survival with a focus on birthweight: A practical breeding perspective. Animal. 2014 Jun 18;Suppl 1:100914.

Heim G, Mellagi APG, Bierhals T, De Souza LP, De Fries HCC, Piuco P, Seidel E, Bernardi ML, Wentz I, Bortolozzo FP. Effects of cross-fostering within 24h after birth on preweaning behaviour, growth performance and survival rate of biological and adopted piglets. Liv Sci. 2012 Dec; 150(1):121-27.

Heub EM, Proll-Cornelissen MJ, Neuhoff C, Tholen E, Grobe-Brinkhaus C. Invited review: Piglet survival: Benefits of immunocompetence. Animal. 2019 Oct;13(10):2114-24.

Howe CG, Claus Henn B, Eckel SP, Farzan SF, Grubbs BH, Chavez TA, Hodes TL, Faham D, Al-Marayati L, Lerner D. Quimby A. Prenatal metal mixtures and birth weight for gestational age in a predominately lower-income Hispanic pregnancy cohort in Los Angeles. Environ Health Perspect. 2020 Nov;128(11):117001.

Huang S, Li N, Jiang L, Dai Z, Li T, Han D, Wang J. Characterization of the early life microbiota development and predominant Lactobacillus species at distinct gut segments of low-and normal-birth-weight piglets. Front Microbiol. 2019 Apr;16(10):797.

Huting AMS, Sakkas P, Wellock I, Almond K, Kyriazakis I. Once small always small? To what extent morphometric characteristics and post-weaning starter regime affect pig lifetime growth performance. Porcine Health Manag. 2018 Jul;23(4):21.

Iida R, Koketsu Y. Climatic factors and risk factors associated with abortion occurrences in Japanese commercial pig herds. Anim Reprod Sci. 2015 Jun;157:78-86.

Johnson JS, Sanz Fernandez MV, Gutierrez NA, Patience JF, Ross JW, Gabler NK, Lucy MC, Safranski TJ, Rhoads RP, Baumgard LH. Effects of in utero heat stress on postnatal body composition in pigs: I. growing phase. J Anim Sci. 2015 Jan;93(1):71-81.

Kaneko M, Iida R, Koketsu Y. Herd management procedures and factors associated with low farrowing rate of female pigs in Japanese commercial herds. Prev Vet Med. 2013 Apr 1;109(1-2):69-75.

Kirkden RD, Broom DM, Andersen IL. Invited review: Piglet mortality: Management solutions. J Anim Sci. 2013 Jul;91(7):3361-89.

Kitkha S, Boonsoongnern A, Ratanavanichrojn N, Jirawattanapong P, Pinyopummin A. Effects of the higher parity and litter size on piglet birth weight and survival rate of later born piglets. Thai J Vet Med. 2017;47(1):79-86.

Knecht D, Srodon S, Duzinski K. The impact of season, parity, and breed on selected reproductive performance parameters of sows. Arch Anim Breed. 2015;58(1):49-56.

Koketsu Y, Iida R. Farm data analysis for lifetime performance components of sows and their predictors in breeding herds. Porcine Health Manag. 2020 Sep;18(6):24.

Koketsu Y, Tani S, Iida R. Factors for improving reproductive performance of sows and herd productivity in commercial breeding herds. Porc Health Manag. 2017 Jan;3(1):1-10.

Langendijk P. Latest advances in sow nutrition during early gestation. Animals (Basel). 2021 Jun 9;11(6):1720.

https://doi.org/10.17221/36/2024-CJAS

Langendijk P, Chen TY, Athorn RZ, Bouwman EG. Embryonic survival at day 9, 21 and 35 of pregnancy in intact and unilaterally oviduct ligated multiparous sows. Anim. 2016 Aug;10(8):1336-41.

Le Dividich J, Charneca R, Thomas F. Relationship between birth order, birth weight, colostrum intake, acquisition of passive immunity and pre-weaning mortality of piglets. Span J Agric Res. 2017 Jun;15(2):e0603.

Li YZ, Gonyou HW. Comparison of management options for sows kept in pens with electronic feeding stations. Can J Anim Sci. 2013 Aug 29;93(1):445-52.

Liu ZH, Zhang XM, Zhou YF, Wang C, Xiong J, Guo LL, Wang L, Jiang SW, Peng J. Effect of increasing feed intake during late gestation on piglet performance at parturition in commercial production enterprises. Anim Reprod Sci. 2020 Jul;(218):106477.

Lopez-Soria S, Moeller SJ, Rakhshandeh A, Lents CA. Effects of birth weight and litter size on pre-weaning piglet mortality, growth, and quality of commercial pigs. J Anim Sci. 2019;97(5):2067-75.

Madsen JG, Bee G. Compensatory growth feeding strategy does not overcome negative effects on growth and carcass composition of low-birth-weight pigs. Anim. 2015 Dec 7;(9):427-36.

Magnabosco D, Cesar E, Cunha P, Bernardi ML, Wentz I, Pandolfo Bortolozzo F. Impact of the birth weight of landrace x large white dam line gilts on mortality, culling and growth performance until selection for breeding herd. Acta Sci Vet. 2015 May;43(1):1274.

Magnabosco D, Bernardi ML, Wentz I, Cunha ECP, Bortolozzo FP. Low birth weight affects lifetime productive performance and longevity of female swine, Liv Sci. 2016 Jan;184(2):119-25.

Marandu N, Halimani TE, Chimonyo M, Shoniwa A, Mutibvu T. Effect of within-litter birth weight variation on piglet survival and pre-weaning weight gain in a commercial herd. J Agric Ext Rural Dev Tropical Subtropical. 2015 Sep;116(2):123–9.

Matabane M, Nephawe K, Thomas R, Maqhashu A, Fhulufhelo R, Netshirovha T, Ngambi J, Tsatsimpe M, Nedambale T. Pre-weaning growth performance of piglets at smallholder farms in Gauteng province. J Agric Sci. 2018 Mar;10(1):18.

Moreira LP, Menegat MB, Barros GP, Bernardi ML, Wentz I, Bortolozzo FP. Effects of colostrum, protein and energy supplementation on survival and performance of low-birth-weight piglets. Liv Sci. 2017 Jun;202(1):188-93.

Morton JM, Langemeier AJ, Rathbun TJ, Davis DL. Immunocrit, colostrum intake, and preweaning body weight gain in piglets after split suckling based on birth weight or birth order. Transl Anim Sci. 2019 Jul 28;3(4):1460-5.

Mulder HA, Rashidi H. Heritable environmental variance causes nonlinear relationship between traits: Application to birth weight and stillbirth of pigs. Genet Res. 2015 Apr;199(4):1255-69.

Muns HS, Ampode KMB, Lagua EB, Park HR, Kim YH, Sharifuzzaman M, Hasan MK, Yang CJ. Backfat thickness at pre-farrowing: Indicators of sow reproductive performance, milk yield, and piglet birth weight in smart farm-based systems. Agriculture. 2023 Dec;14(24):24.

Muns R, Nuntapaitoon M, Tummaruk P. Effect of oral supplementation with different energy boosters in newborn piglets on pre-weaning mortality, growth, and serological levels of IGF-I and IGG1. J Anim Sci. 2017 Jan;95(1):353-60.

Nam NH, Sukon P. Factors influencing within-litter variation of birth weight and the incidence of runt piglets. S Afr J Anim Sci. 2022 Feb;52(1):1-7.

Nam NH, Sukon P. Risk factors associated with stillbirth of piglets born from oxytocin-assisted parturitions. Vet World. 2020 Oct;13(10):2172-7.

Nuntapaitoon M, Muns R, Tummaruk P. Newborn traits associated with e-weaning growth and survival in piglets. Asian-Australas J Anim Sci. 2018 Feb;31(2):237-44.

Oliviero C. Offspring of hyper prolific sows: Immunity, birthweight, and heterogeneous litters. Mol Reprod Dev. 2023 Jul;90(7):580-4.

Pardo C, Berard J, Kreuzer M, Bee G. Intrauterine crowding impairs formation and growth of secondary myofibers in pigs. Animal. 2013 Mar;7(3):430-8.

Pedersen LJ, Larsen MLV, Malmkvist J. The ability of different thermal aids to reduce hypothermia in neonatal piglets. J Anim Sci. 2016, 94(5):2151-59.

Pedersen MLM, Velander IH, Nielsen MBF, Lundeheim N, Nielsen B. Duroc boars have lower progeny mortality and lower fertility than Pietrain boars. Anim Sci. 2019 May 2;3(2):885-92.

Peltoniemi O, Yun J, Bjorkman S, Han T. Coping with large litters: The management of neonatal piglets and sow reproduction. J Anim Sci Technol. 2021 Jan;63(1):1-15.

Pluym LM, Van Nuffel A, Vanweyenberg S, Maes D. Prevalence of lameness and claw lesions during different stages in the reproductive cycle of sows and the impact on reproduction results. Animal. 2013 Jul;7(7):1174-81.

Quesnel H, Quiniou N, Roy H. Supplying dextrose before insemination and L-arginine during the last third of pregnancy in sow diets: effects on withinlitter variation of piglet birth weight. J Anim Sci. 2014 Apr;92(4):1445-50.

Ramirez BC, Hayes MD, Condotta IC, Leonard SM. Impact of housing environment and management on pre-/post-weaning piglet productivity. J Anim Sci. 2022 Jun;100(6):skac142.

https://doi.org/10.17221/36/2024-CJAS

Rasmussen SO, Martin L, Ostergaard MV, Rudloff S, Li Y, Roggenbuck M, Bering SB, Sangild PT. Bovine colostrum improves neonatal growth, digestive function, and gut immunity relative to donor human milk and infant formula in preterm pigs. Am J Physiol Gastrointest Liver Physiol. 2016 Sep 1;311(3):G480-91.

Renteria FJA, Gomez Rosales S, Lopez Hernandez LH, Ordaz Ochoa G, Anaya Escalera AM, Mejia Guadarrama CA, Mariscal Landin G. Main contributions of INIFAP research to swine nutrition in Mexico: Challenges and perspectives. Rev Mex Cienc Pecu. 2021 Nov; 12(1):79-110.

Riddersholm KV, Bahnsen I, Bruun TS, de Knegt LV, Amdi C. Identifying risk factors for low piglet birth weight, high within-litter variation and occurrence of intrauterine growth-restricted piglets in hyperprolific sows. Animals (Basel). 2021 Sep 18;11(9):2731.

Roche R, Shrestha NP, Mekkawy W, Baxter EM, Knap PW, Smurthwaite KM, Jarvis S, Lawrence AB, Edwards SA. Genetic parameters of piglet survival and birth weight from a two-generation crossbreeding experiment under outdoor conditions designed to disentangle direct and maternal effects. J Anim Sci. 2010 Apr;88(4):1276-85.

Roelofs S, Godding L, De Haan JR, Van der Staay FJ, Nordquist RE. Effects of parity and litter size on cortisol measures in commercially housed sows and their offspring. Physiol Behav. 2019 Mar 15;201:83-90.

Romero M, Calvo L, Morales JI, Rodriguez AI, Escudero RM, Olivares A, Lopez-Bote C. Short- and long-term effects of birth weight and neonatal care in pigs. Animals (Basel). 2022 Oct 26;12(21):2936.

Romero M, Calvo L, Morales JI, Magro A, Rodriguez AI, Segura J, Escudero R, Lopez-Bote C, Olivares A. Short- and long-term effects of split-suckling in pigs according to birth weight. Animals (Basel). 2023 Nov 14;13(22):3521.

Rothschild MF, Ruvinsky A. The genetics of the pig. 2nd ed. Wallington, UK: CAB International; 2011. p. 218-41,325-54,390-425.

Rutherford KMD, Baxter EM, D'eath RB, Turner SP, Arnott G, Roehe R, Ask B, Sandoe P, Moustsen VA, Thorup F, Edwards SA, Berg P, Lawrence AB. The welfare implications of large litter size in the domestic pig I: Biological factors. Anim Welf. 2013 May;22:199-218.

Sangild PT, Vonderohe C, Melendez Hebib V, Burrin DG. Potential benefits of bovine colostrum in pediatric nutrition and health. Nutrients. 2021 Jul 26;13(8):2551.

Schinckel AP, Elinstein ME, Stewart TS, Schwab C, Olynkf NJ. Use of a stochastic model to evaluate the growth performance and profitability of pigd from different litter sizes and parities of dams. Pr Anim Sci. 2010 Oct;26(5):547-60.

Schmitt O, Baxter EM, Boyle LA, O Driscoll K. Nurse sow strategies in the domestic pig: II. Consequences for piglet growth, suckling behavior, and sow nursing behavior. Animal. 2019 Mar;13(3):590-9.

Serenius T, Stalder KJ, Karriker LA. Pre-weaning mortality in pigs. Vet Clin N Am – Food Anim Pract. 2019; 35(3):515-28.

Silva EG, dos SO, Rangel AH do N, Murmam L, Bezerra MF, Oliveira JPF de. Bovine colostrum: Benefits of its use in human food. Food Sci Technol. 2019 Dec;39(1):355-62.

Soede NM, Kemp B. Best practices in the lactating and weaned sow to optimize reproductive physiology and performance. In: Farmer C, editor. The gestating and lactating sow. Wageningen, Wageningen Academic; 2015. p. 99-106.

Swiatkiewicz M, Hanczakowska E, Okon K, Kowalczyk P, Grela ER. Effect of maternal diet and medium chain fatty acids supplementation for piglets on their digestive tract development, structure, and chyme acidity as well as performance and health status. Animals (Basel). 2020 May 11;10(5):834.

Swinbourne AM, Kelly JM, Kind KL, Kennaway DJ, Van Wettere WHEJ. The effects of season and moderate nutritional restriction on ovarian function and oocyte nuclear maturation in cycling gilts. Theriogenology. 2014 Dec;82(9):1303-9.

Tang T, Gerrits WJ, Soede NM, van der Peet-Schwering CM, Reimert I. Effects of timing of grouping and split weaning on growth performance and behaviour of piglets in a multi-suckling system. Appl Anim Behav Sci. 2023 Feb;259(1):105835.

Theil PK, Lauridsen C, Quesnel H. Neonatal piglet survival. Impact of sow nutrition around parturition on fatal glycogen deposition and production and composition of colostrum and transient milk. Animal. 2014 Jul;8(7):1021-30.

Tian Z, Wang LG, Shi HB, Hua Y, Zhang LC, Xin L, Lei P, Jing L, Zhang YB, Zhao KB, Li-Xian W. Heritability and genetic and phenotypic correlations of litter uniformity and litter size in large white sows. J Integr Agric. 2016 Apr;15(1):848-54.

Tucker BS, Craig JR, Morrison RS, Smits RJ, Kirkwood RN. Piglet viability: A review of identification and pre-weaning management strategies. Animals (Basel). 2021 Oct 6; 11(10):2902.

Tummaruk P, Sang-Gassanee K. Effect of farrowing duration, parity number and the type of anti-inflammatory drug on post-parturient disorders in sows: A clinical study. Trop Anim Health Prod. 2013 Apr;45(4):1071-7.

Vallet JL, Stalder KJ, Phillips C, Cushman RA. Litter-of-origin trait effects on gilt development. J Anim Sci. 2016 Jan;94(1):96-105.

https://doi.org/10.17221/36/2024-CJAS

Van den Bosch M, Wijnen J, van de Linde IB, van Wesel AA, Melchior D, Kemp B, Clouard C, van den Brand H. Effects of maternal dietary nitrate supplementation during the perinatal period on piglet survival, body weight, and litter uniformity. Transl Anim Sci. 2019 Feb 7;3(1):464-72.

Vandaele M, Van Kerschaver C, Degroote J, Van Ginneken C, Michiels J. Piglet performance and colostrum intake in litters either or not split-suckled during the first day or during the first three days of life. Liv Sci. 2020 Nov;241(1):104265.

Vande Pol KD, Bautista RO, Olivo A, Harper H, Shull CM, Brown CB, Ellis M. Effect of rearing cross-fostered piglets in litters of differing size relative to sow functional teat number on preweaning growth and mortality. Transl Anim Sci. 2021 Oct 2;5(4):txab193.

VanderWaal K, Deen J. Global trends in infectious diseases of swine. Proc Natl Acad Sci. 2018; Nov 6;115(45):11495-500.

Vodolazska D, Feyera T, Lauridsen C. The impact of birth weight, birth order, birth asphyxia, and colostrum intake per se on growth and immunity of the suckling piglets. Sci Rep. 2023 May 17;13(1):8057.

Vodolazska D, Lauridsen C. Effects of dietary hemp seed oil to sow on fatty acid profiles, nutritional and immune status of piglets. J Anim Sci Biotechnol. 2020 Mar 18;11:28.

Ward SA, Kirkwood RN, Plush KJ. Are larger litters a concern for piglet survival or an effectively manageable trait? Animals (Basel). 2020 Feb 17;10(2):309.

Wellock IJ, Emmans GC, Kyriazakis I, Houdijk JG. The consequences of genetic selection for efficiency in livestock: The paradox of reduced animal welfare. Anim. 2014;8(1):23-32.

Wen B, Njunge JM, Bourdon C, Gonzales GB, Gichuki BM, Lee D, Wishart DS, Ngari M, Chimwezi E, Thitiri J, Mwalekwa L, Voskuijl W, Berkley JA, Bandsma RH. Systemic inflammation and metabolic disturbances underlie inpatient mortality among ill children with severe malnutrition. Sci Adv. 2022 Feb 18;8(7):eabj6779.

Wientjes JGM, Soede NM, Knol E, Van den Brand H, Kemp B. Piglet birth weight and litter uniformity: Effects of weaning-to-pregnancy interval and body condition changes in sows of different parities and crossbred lines. J Anim Sci. 2013 May;91(5):2099-107.

Yuan TL, Zhu YH, Shi M, Li TT, Li N, Wu GY, Wang JJ. Within-litter variation in birth weight: Impact of nutritional status in the sow. J Zhejiang Univ Sci B. 2015 Jun;16(6):417-35.

Zeng ZK, Urriola PE, Dunkelberger JR, Eggert JM, Vogelzang R, Shurson GC, Johnston LJ. Implications of early-life indicators for survival rate, subsequent growth performance, and carcass characteristics of commercial pigs. J Anim Sci. 2019 Jul 30;97(8):3313-25.

Zhang X, Wang M, He T, Long S, Guo Y, Chen Z. Effect of different cross-fostering strategies on growth performance, stress status and immunoglobulin of piglets. Animals (Basel). 2021 Feb 14;11(2):499.

Zindove TJ, Dzomba EF, Kanengoni AT, Chimonyo M. Effects of within-litter birth weight variation of piglets on performance at 3 weeks of age and at weaning in a Large White x Landrace sow herd. Liv Sci. 2013 Aug;155(1):348-54.

Zotti E, Resmini FA, Schutz LG, Volz N, Milani RP, Bridi AM, Alfieri AA, Silva CAD. Impact of piglet birthweight and sow parity on mortality rates, growth performance, and carcass traits in pigs. Animal. 2013 Nov;7(11):1884-92.

Received: March 14, 2024
Accepted: July 15, 2024