

The use of indirect methods for the prediction of lucerne quality in the first cut under the conditions of Central Europe

J. HAKL, J. ŠANTRŮČEK, P. FUKSA, L. KRAJÍČ

Department of Forage Crops and Grassland Management, Czech University of Life Sciences, Prague, Czech Republic

ABSTRACT: The goal of this study was to compare the growing degree-days (GDD) and predictive equations for alfalfa quality (PEAQ) for the prediction of lucerne quality and to test their accuracy and suitability in the first cut period in Central Bohemia. Several additional stand parameters were verified in order to increase the accuracy of the quality estimate under these environmental conditions. In 2004–2007, the measurement and sampling were repeatedly realized from the late vegetative to the early bloom stage in six replications. For the GDD model across the years, the obtained R^2 for NDF, ADF and CP were 0.40, 0.57 and 0.65, respectively. It seems that the forage quality response to accumulated GDD was related to the stand development and it could be the reason for low R^2 across all the years. For the PEAQ model, R^2 were 0.62, 0.92, and 0.85, respectively. Similarly like in the GDD model, the effect of stand development across the years on changes in the slopes of equations was observed. The accuracy of the model combination was not higher in comparison with the PEAQ model. The count of stems per plant, density of stems per m^2 , dry matter yield and average stem weight improved the NDF content prediction within a four-year period. In these models, variables which represent the stand development should be taken into account.

Keywords: forage; alfalfa; quality; stand; fibre

The timing of spring forage harvest is critical for obtaining optimal quality for animal production. For forage that serves as the primary fibre source in the diet, NDF is the principal forage quality variable of concern (Parsons et al., 2006a). Some predictive equations can be used to estimate the forage quality of lucerne, assisting the producers in decision making at harvest time. Parsons et al. (2006b) described an ideal method for estimating lucerne quality in the field as a harvest decision aid must be quick, simple, inexpensive, and consistent across all harvests during the season and across a wide range of environments.

A number of methods have been developed to estimate lucerne NDF, including models based on weather, chronological age and plant morphology (Fick et al., 1994). These parameters are related not only to NDF or ADF but also to CP, DOM, and IVDDM (Sulc et al., 1997; Rinne and Nykanen, 2000; Mitchell et al., 2001). Near infrared reflectance spectroscopy (NIRS) belongs to indirect methods (Mika et al., 2003) but this method is not so suitable for field conditions, moreover, special instrumentation is needed.

The most widely used of these methods are predictive equations for alfalfa quality (PEAQ). This

method is based on the length of the tallest stem and the stage of the most mature stem in the sample (Hintz and Albrecht, 1991). The equations also performed well across the spring and summer growth cycles (Sulc et al., 1997) although lucerne quality varied among growth cycles during the year (Niwinska et al., 2005), which is connected with changes in the leaf stem ratio (Tyrolová and Výborná, 2008). Similarly Skládanka et al. (2008) documented changes in fibre content in forage of grasses at the end of the growing season. These equations have been developed for many regions of the USA. Results indicated some bias when the equations were used outside the state of development; however, the prediction errors were sufficiently low to suggest the PEAQ equations are robust over a wide range of environments (Parsons et al., 2006a). Parsons et al. (2006b) described some modified equations, based only on lucerne height.

Growing degree-days (GDD) are a temperature-derived index representing the amount of heat to which plants are exposed. This method has been used successfully in maize development, but has had a mixed success in the perennial types of forage (Sulc et al., 1999). The GDD for lucerne are calculated as follows. Average the maximum and minimum temperature for each day (24 h period) beginning March 1, subtract the base temperature ($41^{\circ}\text{F} = 5^{\circ}\text{C}$) and sum up the growing degrees for all days that have a positive number (Allen and Beck, 1996). The GDD are not reliable when the soil moisture is inadequate, which often occurs during the second cutting growth. It is so presumably because

forage growth is limited more by soil moisture than by heat units. Neither GDD nor PEAQ adequately predicted NDF for the third cutting lucerne (Lee et al., 2002). Similarly, the chronological age of forage relates to quality in spring growth, but it is inconsistently related to quality in regrowth (Sulc et al., 1999). In Europe, the GDD method has proved to be an accurate predictor of grass digestibility under Finnish conditions (Rinne and Nykanen, 2000). The base temperature is 5°C as published by Allen and Beck (1996).

The above discussed methods are quick, easy and their accuracy is acceptable under the USA conditions (Owens et al., 1995; Allen and Beck, 1996; Sulc et al., 1997). Since no single method will ever result in a perfect prediction of lucerne quality in the future, a combination of methods may be most acceptable (Sulc et al., 1999). Allen and Beck (1996) suggested that GDD in combination with the plant height and the maturity stage might be more accurate for estimating lucerne NDF than using GDD alone. Some additional variables for improving accuracy in models were used by Parsons et al. (2006a) in a lucerne grass mixture. As noted by Hakl et al. (2006), the stand structure could influence the fibre content of lucerne but this effect is low in dense stand. It is important to understand that any method used to predict or estimate forage quality will have an error in relation to wet chemistry analyses (Sulc et al., 1999).

In the Czech Republic, these methods have not yet been tested for any perennial forage crops. The goal of this study was to assess preliminary equations of these methods and to test their accuracy and suitability for

Table 1. Descriptions and ranges of variables evaluated as potential predictors of lucerne quality in lucerne stand

Variables	Description	Min.	Max.
ASW	average stem weight (g)	2	21
D	stand density (plants/m ²)	75	700
DMY	dry matter yield (kg/ha)	2 000	17 775
GDD	accumulated growing degree days ($^{\circ}\text{C}$), base 5°C	193	578
MSL	height of the tallest lucerne stems in the sample area (cm)	29.2	143.0
S	morphological stage of development of the most mature lucerne stems in the sample area	2	5
SPP	count of stems per plant (pcs/plant)	2.2	15.3
TSC	total stem count (pcs/m ²)	375	2 375
Y	year of vegetation	2	5

Table 2. Lucerne developmental stage used to assign a numerical value to the most mature stems in the sampling area, codes of stages and descriptions are according to Kalu and Fick (1981)

Developmental stages	Code	Description
Late vegetative	2	stems length > 300 mm, no visible buds
Early bud	3	1 to 2 nodes with visible buds
Late bud	4	> 2 nodes with visible buds
Early bloom	5	1 node with at least 1 open flower

lucerne in the first cut period in Central Bohemia. In these models, we also verified some additional stand parameters for increasing the accuracy of the quality estimate in our environmental conditions.

MATERIAL AND METHODS

In spring 2003, the lucerne experiment was established at the Czech University of Life Sciences in Prague in Central Bohemia (286 m above sea level, 50°08' N, 14°24' E). The long-term annual temperature is 9.3°C and precipitation 510 mm.

The experimental plants (*Medicago sativa* L.) consisted of two lucerne candivars (ŽE XLII, ŽE XLV) and the Jarka variety grown in two replications on completely randomised blocks. In 2004–2007, the measurements and samplings were repeatedly realized from the late vegetative to the early bloom stage in the first cut per each treatment in a 20 × 20 cm square. Stand density, total stem count, dry matter yield, and count of stems per plant were assessed for each sample. The height of the tallest lucerne stem in the sample area was measured to the terminal point of the stem. Abbreviations of the variables, their descriptions and ranges are summa-

Table 3. Means of yield and stand structure parameters

Year	S	D (plants/m ²)		TSC (pcs/m ²)		MSL (cm)		SPP (pcs/plant)		ASW (g)	
		mean	SE	mean	SE	mean	SE	mean	SE	mean	SE
2004	2	412	56	1 254	166	33.0	1.5	3.07	0.17	3.05	0.42
	3	433	68	1 162	170	48.9	2.3	2.73	0.10	3.59	0.37
	4	454	50	1 437	146	63.0	4.2	3.21	0.26	5.55	0.86
	5	354	50	1 037	90	73.8	4.1	3.05	0.24	7.48	0.28
2005	2	337	37	1 537	150	45.3	1.7	4.60	0.24	3.69	0.46
	3	204	16	979	93	60.3	2.3	4.81	0.31	5.89	0.53
	4	279	18	970	135	74.1	3.0	3.47	0.45	8.68	0.86
	5	258	11	941	58	106.1	3.2	3.70	0.32	12.33	1.13
2006	2	154	12	1 033	295	60.8	1.8	6.62	1.64	6.24	1.28
	3	187	11	1 341	106	81.1	1.4	7.19	0.50	7.62	0.78
	4	192	18	1 088	66	98.8	5.5	5.85	0.48	11.62	0.94
	5	150	18	800	100	120.3	6.1	5.34	0.27	14.83	1.70
2007	2	117	14	933	129	52.8	2.1	8.23	1.12	6.02	0.51
	3	125	17	1 133	44	70.8	3.4	9.96	1.45	8.73	0.99
	4	121	10	913	70	79.3	1.8	7.81	0.94	10.44	1.05
	5	117	17	838	91	94.8	2.6	7.60	0.87	14.42	1.65

SE = standard error; $n = 6$; for the description of variables see Table 1

Table 4. Means of forage yield and quality

Year	S	Date	GDD (°C)	DMY (kg/ha)		CP (%)		ADF (%)		NDF (%)	
				mean	SE	mean	SE	mean	SE	mean	SE
2004	2	12.5.	324	3 516	207	24.67	0.39	25.41	0.99	30.21	0.66
	3	21.5.	405	3 879	417	22.58	0.53	29.69	0.77	34.46	0.96
	4	31.5.	471	7 550	729	18.94	0.54	33.66	1.06	37.12	1.32
	5	9.6.	578	7 770	764	20.28	0.71	34.86	1.30	40.36	1.58
2005	2	4.5.	262	5 512	744	24.23	0.99	26.07	4.59	40.72	2.43
	3	19.5.	341	5 608	410	23.50	0.40	30.00	1.05	45.38	1.02
	4	25.5.	389	8 108	1 052	21.40	0.61	32.93	0.66	43.10	1.81
	5	7.6.	567	11 512	1 184	16.57	0.81	43.87	1.15	52.87	1.25
2006	2	10.5.	265	5 400	1 106	26.11	0.84	29.82	1.28	34.72	0.76
	3	16.5.	324	10 279	1 415	23.78	0.47	32.70	1.07	36.85	0.76
	4	25.5.	416	12 616	1 269	20.63	0.43	36.92	0.52	43.17	0.51
	5	9.6.	512	11 895	1 817	16.47	0.80	41.62	2.01	47.68	1.80
2007	2	24.4.	193	5 345	544	25.12	0.81	26.13	0.95	30.08	1.28
	3	2.5.	258	9 775	951	21.55	0.42	31.77	0.86	35.67	0.68
	4	7.5.	300	9 663	1 433	21.23	0.57	35.28	1.77	40.12	2.41
	5	16.5.	394	11 446	684	19.16	0.55	35.32	1.07	43.77	1.68

SE = standard error; $n = 6$; for the description of variables see Table 1

rized in Table 1. The lucerne maturity categories of Kalu and Fick (1981) (Table 2) were used to assign a numerical value to the most mature stem in the sample area. All samples were oven-dried at 60°C, homogenised to a particle size of 1 mm, and analysed for crude protein (CP), neutral detergent fibre (NDF), and acid detergent fibre (ADF) content. CP was quantified using the Kjeldahl procedure for N determination ($\% \text{ CP} = \% \text{ N} \times 6.25$; AOAC, 1990). The NDF and ADF contents were determined according to Van Soest et al. (1991). GDD for each harvest date were calculated according to Allen and Beck (1996). Direct and stepwise linear regression analyses were performed using Statistica 6.1 (StatSoft, 2003).

RESULTS AND DISCUSSION

The means and standard errors of measured parameters are shown in Tables 3 and 4. You can see increasing DMY and MSL during the follow up first cut periods. Within a four-year period, increas-

ing SPP was caused by a natural decrease of stand density and subsequent plant development which is associated with changes in the root traits (Hakl et al., 2007). The changes in lucerne quality during the first cut periods correspond with changes described by Homolka et al. (2008). No significant differences among lucerne entries in forage quality were detected, therefore this factor was not included in subsequent analyses.

As noted by Lee et al. (2002), the GDD is not reliable when the soil moisture is inadequate. During the first cut period, this site did not show any signs of moisture deficit in all experimental years. The models of the GDD method are shown in Table 5. The R^2 and standard error of prediction (SEP) values varied from one year to the next. This is in accordance with Allen and Beck (1996), who reported that prediction equations developed in one year were not stable across the years. Cherney (1995) found that GDD prediction equations of NDF content developed for different data sets had considerably different intercepts, but the slopes of the equations were relatively consistent. In our ex-

Table 5. Relating lucerne quality (y ; NDF, ADF, CP; %) and accumulated growing degree-days (x ; GDD; °C) in the first cut period (linear regression, $y = a + bx$)

Y quality	Year	X = GDD model	SEP	R^2	P -value	n
NDF	2004	$18.06 + 0.039 \text{ GDD}$	2.81	0.65	0.0000	24
	2005	$30.59 + 0.038 \text{ GDD}$	4.33	0.52	0.0000	24
	2006	$19.91 + 0.055 \text{ GDD}$	2.58	0.81	0.0000	24
	2007	$17.85 + 0.068 \text{ GDD}$	4.01	0.63	0.0000	24
	2004–2007	$26.77 + 0.035 \text{ GDD}$	4.96	0.40	0.0083	16*
ADF	2004	$14.06 + 0.038 \text{ GDD}$	2.72	0.65	0.0000	24
	2005	$10.16 + 0.059 \text{ GDD}$	5.73	0.59	0.0000	24
	2006	$17.23 + 0.048 \text{ GDD}$	3.11	0.69	0.0000	24
	2007	$19.27 + 0.045 \text{ GDD}$	3.42	0.50	0.0001	24
	2004–2007	$19.90 + 0.035 \text{ GDD}$	3.54	0.57	0.0008	16*
CP	2004	$30.02 - 0.019 \text{ GDD}$	1.88	0.49	0.0001	24
	2005	$31.79 - 0.026 \text{ GDD}$	1.79	0.75	0.0000	24
	2006	$36.37 - 0.039 \text{ GDD}$	1.55	0.85	0.0000	24
	2007	$29.77 - 0.028 \text{ GDD}$	1.57	0.65	0.0000	24
	2004–2007	$29.40 - 0.021 \text{ GDD}$	1.77	0.65	0.0001	16*

*for elimination of heterogeneity on sampling days the average values from Table 3 and 4 were used

SEP = standard error of prediction, intercepts (a) and slopes (b) significant at $P < 0.05$

Table 6. Relating lucerne quality (y ; NDF, ADF, CP; %) and stage of development (x_1 ; S; code 2–5) and maximal stem length (x_2 , MSL; cm) in the first cut period (PEAQ method, linear regression, $y = a + b_1x_1 + b_2x_2$)

Y quality	Year	X model	SEP	R^2	P -value	n
NDF	2004	$21.95 - 0.07 \text{ S} + 0.30 \text{ MSL}$	1.56	0.90	0.0000	24
	2005	$32.95 - 0.99 \text{ S} + 0.22 \text{ MSL}$	4.54	0.49	0.0008	24
	2006	$23.83 + 3.67 \text{ S} + 0.04 \text{ MSL}$	2.72	0.80	0.0000	24
	2007	$26.58 + 7.06 \text{ S} - 0.19 \text{ MSL}$	3.77	0.69	0.0000	24
	2004–2007	$24.11 + 1.85 \text{ S} + 0.13 \text{ MSL}$	4.11	0.62	0.0019	16*
ADF	2004	$17.99 + 0.05 \text{ S} + 0.23 \text{ MSL}$	1.92	0.83	0.0000	24
	2005	$12.86 + 1.05 \text{ S} + 0.23 \text{ MSL}$	5.95	0.58	0.0001	24
	2006	$22.14 + 4.64 \text{ S} - 0.03 \text{ MSL}$	3.20	0.69	0.0000	24
	2007	$25.30 + 5.10 \text{ S} - 0.15 \text{ MSL}$	3.14	0.60	0.0000	24
	2004–2007	$16.89 + 1.67 \text{ S} + 0.14 \text{ MSL}$	1.61	0.92	0.0000	16*
CP	2004	$27.75 - 1.17 \text{ S} - 0.04 \text{ MSL}$	1.77	0.57	0.0000	24
	2005	$30.54 - 0.63 \text{ S} - 0.10 \text{ MSL}$	1.92	0.72	0.0000	24
	2006	$33.30 - 2.90 \text{ S} - 0.02 \text{ MSL}$	1.65	0.84	0.0000	24
	2007	$27.20 - 2.28 \text{ S} + 0.03 \text{ MSL}$	1.60	0.65	0.0000	24
	2004–2007	$29.99 - 2.01 \text{ S} - 0.02 \text{ MSL}$	1.20	0.85	0.0000	16*

*for elimination of heterogeneity on sampling days the average values from Table 3 and 4 were used

SEP = standard error of prediction, intercepts (a) and slopes (b_1 , b_2) significant at $P < 0.05$

periment, the intercepts of equations for the NDF prediction between years were relatively consistent, except for the year 2005. The slopes increased from 0.039 in 2004 to 0.068 in 2007. In 2004, the lowest slopes were also recorded for ADF and CP. We can assume that the utilization of heat units for fibre production varied in dependence on the stand structure when an older stand with more developed plants increased the fibre content faster than a younger stand with less developed plants. This effect was most obvious in the case of NDF with continually increasing slopes from 2004 to 2007. It is possible to conclude that the forage quality response to accumulated GDD was related to stand development in the four-year period. It could be the reason for low R^2 and high SEP values of prediction equations across all the years. In this case, the best estimate was recorded for CP where the R^2 value was 0.65.

The models of PEAQ are shown in Table 6. We used the 4-stage scale according to Table 2. In contrast to GDD models, the slopes in equations of each year were not always significantly different from zero. Similarly like in the GDD model, the R^2 and SEP values varied from one year to the next. The prediction equations across all the years were more accurate than the GDD models. The original equations according to Hintz and Albrecht (1991) reached the R^2 values of 0.89 for NDF ($= 16.89 + 0.81 S + 0.27 \text{ MSL}$), 0.88 for ADF ($= 11.57 + 0.79 S + 0.21 \text{ MSL}$) and 0.74 for CP ($= 30.71 - 0.89 S - 0.09 \text{ MSL}$). Sulc et al. (1997)

evaluated the performance of PEAQ across five states and presented the R^2 value 0.74 for NDF ($= 23.67 + 0.41 S + 0.21 \text{ MSL}$) and 0.73 for ADF prediction. According to Owens et al. (1995), the CP was not predicted accurately using PEAQ. In the first cut, they obtained the R^2 values of 0.83 for NDF ($= 18.19 + 1.44 S + 0.21 \text{ MSL}$), 0.78 for ADF ($= 11.92 + 1.11 S + 0.18 \text{ MSL}$) and 0.37 for CP ($= 29.63 - 0.87 S - 0.09 \text{ MSL}$). In our experiment, the R^2 values were 0.62, 0.92, and 0.85, respectively. Similarly like in the GDD model, the effect of stand development across years on changes in slopes of the equations was observed. This effect was most obvious in the case of fibre with increasing slopes of maturity and decreasing slopes of MSL from 2004 to 2007 with opposite trends for CP. Generally, it seems that across years the slopes for stage were higher and for MSL they were lower in comparison with the equations mentioned above.

The combinations of models are shown in Table 7. The combination of GDD and PEAQ models was better for all quality parameters than GDD alone, which is in accordance with suggestions of Sulc et al. (1999) that the combination of methods might be most acceptable. The accuracy of the model combination was not higher in comparison with the PEAQ model. The highest accuracy was observed for ADF, the lowest for NDF. The GDD model alone is less accurate than PEAQ but, according to Sulc et al. (1999), it eliminates the need to sample a field, and therefore eliminates the potential for lucerne sampling error.

Table 7. Combination of models and stepwise regression for the prediction of lucerne quality (NDF, ADF, CP; %); linear regression

	Model*	SEP	R^2	P-value
Combination of GDD and PEAQ				
NDF	$22.46 + 0.15 \text{ MSL} + 0.259 S + 0.015 \text{ GDD}$	4.16	0.64	0.0054
ADF	$15.28 + 0.16 \text{ MSL} + 0.117 S + 0.014 \text{ GDD}$	1.37	0.94	0.0000
CP	$30.61 - 0.03 \text{ MSL} - 1.41 S - 0.006 \text{ GDD}$	1.19	0.86	0.0000
Stepwise regression				
NDF	$48.01 + 0.017 \text{ TSC} - 3.72 \text{ SPP} - 0.08 \text{ DMY} + 2.95 S + 2.87 \text{ ASW}$	3.20	0.82	0.0017
ADF	$15.24 + 0.16 \text{ MSL} + 0.015 \text{ GDD}$	1.32	0.94	0.0000
CP	$29.72 - 2.31 S$	1.18	0.84	0.0000

*for the description of model variables see Table 1

SEP = standard error of prediction, intercepts and slopes significant at $P < 0.05$, $n = 16$

From other variables, the stepwise algorithm included SPP, DMY, TSC and ASW to models for NDF prediction whilst ADF was the best for the model using MSL and GDD. CP prediction was the best using only the stage of development (Table 7). The accuracy of stepwise models was similar to the PEAQ model except for NDF where the R^2 value increased to 0.82. The significant slopes of selected variables indicate their importance for fibre prediction. The SPP variable is a function of D and TSC. It corresponds with Šantrůček (1989), who stated that stand density can influence the fibre content although this effect is low in a dense stand (Hakl et al., 2006). In our experiment, these variables represent a level of stand development where they correspond with the changes of stand in a four-year period. It seems that the accuracy of the models could be influenced by these changes. The inclusion of variables which characterized the stand development in the models can improve the accuracy of the fibre estimate and thus it should be taken into account.

CONCLUSIONS

The PEAQ method seems to be a promising method for the lucerne quality prediction under the conditions in Bohemia. The GDD method was less accurate across the years and the combination with the PEAQ did not improve the quality of the models. However, the GDD method eliminates the need to sample a field, so it would be suitable as the signal method for starting with field evaluation. In both methods, the accuracy of estimates was the lowest for NDF prediction. According to our opinion, the combination of developmental stage and stand height or GDD seems to be a promising method for predicting harvest dates in the Czech Republic and thus suitable for field experiments as well as for farm use. To develop a functional model, experiments on many sites are needed with a follow up validation of this model. In our opinion, the variables which represent stand development should be taken into account in these models mainly for NDF prediction.

REFERENCES

- Allen M., Beck J. (1996): Relationship between spring harvest alfalfa quality and growing degree days. In: Proc. 26th Natl. Alfalfa Symposium, East Lansing. CASC, Davis, USA, 16–25.
- AOAC (1990): Official methods of analysis. 15th ed. AOAC, Arlington, USA.
- Cherney J.H. (1995): Spring alfalfa harvest in relation to growing degree days. In: 25th Proc. Natl. Alfalfa Symposium. Syracuse, New York, Woodland, USA, 29–36.
- Fick G.W., Wilkens P.W., Cherney J.H. (1994): Modeling forage quality changes in the growing crop. In: Fahey G.C. Jr. (ed.): Forage Quality, Evaluation, and Utilization. ASA, CSSA, and SSSA, Madison, USA, 757–795.
- Hakl J., Šantrůček J., Kalista J., Kocourková D. (2006): The influence of lucerne stand structure on fibre content in the forage. In: Proc. 21st EGF Meeting. Badajoz, Spain, 264–266.
- Hakl J., Šantrůček J., Kalista J., Kocourková D. (2007): The root morphology of new Czech lucerne cultivars. *Scientia Agriculturae Bohemica*, 38, 1–5.
- Hintz R.W., Albrecht K.A. (1991): Prediction of alfalfa chemical composition from maturity and plant morphology. *Crop Science*, 31, 1561–1565.
- Homolka P., Koukolová V., Němec Z., Mudřík Z., Hučko B., Sales J. (2008): Amino acid contents and intestinal digestibility of lucerne in ruminants as influenced by growth stage. *Czech Journal of Animal Science*, 53, 499–505.
- Kalu B.A., Fick G.W. (1981): Quantifying morphological development of alfalfa for studies of herbage quality. *Crop Science*, 21, 267–271.
- Lee K., Allen M., Leep R. (2002): Predicting optimum time of alfalfa harvest. In: Proc. Tri-state Dairy Nutrition Conference, Fort Wayne. Ohio State University, Columbus, USA, 149–152.
- Mika V., Pozdíšek J., Tillmann P., Nerušil P., Buchgraber K., Gruber L. (2003): Development of NIR calibration valid for two different grass sample collections. *Czech Journal of Animal Science*, 48, 419–424.
- Mitchell R., Fritz J., Moore K., Moser L., Vogel K., Redfearn D., Wester D. (2001): Predicting forage quality in Switchgrass and Big Bluestem. *Agronomy Journal*, 93, 118–124.
- Niwinska B., Strzetelski J.A., Kowalczyk J., Borowiec F., Domanski P. (2005): The effect of phenological stage and season on nutritive value, chemical composition and nutrient digestibility of lucerne (*Medicago sativa* L.) green forage in the alimentary tract of cattle. *Czech Journal of Animal Science*, 50, 511–518.
- Owens V.N., Albrecht K.A., Hintz R.W. (1995): A rapid method for predicting alfalfa quality in the field. *Journal of Production Agriculture*, 8, 491–495.

- Parsons D., Cherney J.H., Gauch H.G. (2006a): Estimation of preharvest fiber content of mixed alfalfa-grass stand in New York. *Agronomy Journal*, 98, 1081–1089.
- Parsons D., Cherney J.H., Gauch H.G. (2006b): Alfalfa fiber estimation in mixed stands and its relationship to plant morphology. *Crop Science*, 46, 2446–2452.
- Rinne M., Nykanen A. (2000): Timing of primary growth harvest affects the yield and nutritive value of timothy-red clover mixtures. *Agricultural and Food Science in Finland*, 9, 121–134.
- Šantrůček J. (1989): Quality of lucerne forage in dependence on soil compacting and cultivation. *Rostlinná výroba*, 35, 1101–1107.
- Skládanka J., Dohnal V., Ježková A. (2008): Fibre and ergosterol contents in forage of *Arrhenatherum elatius*, *Dactylis glomerata* and *Festulolium* at the end of the growing season. *Czech Journal of Animal Science*, 53, 320–329.
- StatSoft Inc. (2003): *Statistica for Windows*. StatSoft, Tulsa, USA.
- Sulc R.M., Albrecht K.A., Cherney J.H., Hall M.H., Mueller S.C., Orloff B. (1997): Field testing of rapid method for estimating alfalfa quality. *Agronomy Journal*, 89, 952–957.
- Sulc R.M., Albrecht K.A., Owens V.N., Cherney J.H. (1999): Update on predicting harvest time for alfalfa. In: *Proc. Tri-state Dairy Nutrition Conference*. Fort Wayne, Ohio State University, Columbus, USA, 167–177.
- Tyrolová Y., Výborná A. (2008): Effect of the stage of maturity on the leaf percentage of lucerne and the effect of additives on silage characteristic. *Czech Journal of Animal Science*, 53, 330–335.
- Van Soest P.J., Robertson J.B., Lewis B.A. (1991): Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74, 3583–3597.

Received: 2009–04–08

Accepted after corrections: 2009–12–14

Corresponding Author

Ing. Josef Hakl, Ph.D., Department of Forage Crops and Grassland Management, Czech University of Life Sciences, Prague, Kamýcká 129, 165 21 Prague 6-Suchbát, Czech Republic
Tel. +420 224 383 038, fax +420 234 381 831, e-mail: hakl@af.czu.cz

INSTITUTE OF AGRICULTURAL ECONOMICS AND INFORMATION

Mánesova 75, 120 56 Prague 2, Czech Republic

Tel.: + 420 222 000 111, Fax: + 420 227 010 116, E-mail: redakce@uzei.cz

Account No. 86335-011/0100 KB

IBAN – CZ220100000000086335011; SWIFT address – KOMBCZPPXXX

In this institute scientific journals dealing with the problems of agriculture and related sciences are published on behalf of the Czech Academy of Agricultural Sciences. The periodicals are published in English.

Journal	Number of issues per year	Yearly subscription in USD
Plant, Soil and Environment	12	540
Czech Journal of Animal Science	12	660
Agricultural Economics (Zemědělská ekonomika)	12	540
Journal of Forest Science	12	480
Veterinární medicína (Veterinary Medicine – Czech)	12	720
Czech Journal of Food Sciences	6	420
Plant Protection Science	4	140
Czech Journal of Genetics and Plant Breeding	4	160
Horticultural Science	4	160
Research in Agricultural Engineering	4	140
Soil and Water Research	4	140

Subscription to these journals be sent to the above-mentioned address.

INSTRUCTIONS TO AUTHORS

Czech Journal of Animal Science

The journal publishes original scientific papers and board-invited review articles in English. The author is fully responsible for the originality of the paper and its subject and formal correctness. The author's declaration that the paper has not been published anywhere else should be enclosed. The Board of Editors decides on the publication of papers, taking into account peer reviews, scientific importance, and manuscript quality. Good laboratory practice and ethical rules must be followed. The SI international system of measurement units should be used. Manuscripts must be grammatically and linguistically correct and authors whose native language is not English are advised to seek the help of native English-speaker. Manuscript containing language errors are prejudiced in the reviewing process and may be returned to the author for rewriting before peer review and/or before acceptance.

Only manuscripts assessed by leading experts in the field will be published. If such reviews are not available within four months after registration of the manuscript, the peer review process is ceased, and the authors are notified. They can resubmit the manuscript, after its thorough revision and/or update, either to Czech Journal of Animal Science for a new assessment or to another journal. This should eliminate a long waiting period, and likely rejection of the manuscript.

If a revision of manuscript following the recommendation of the reviewers is requested, the modified manuscript must be re-submitted within six weeks. The authors can, however request an extension of the re-submission deadline, if necessary. All parts of the manuscript, including tables and figures (even unchanged) must be re-submitted. A detailed reply by the authors to every point of reviewer's recommendations must be attached to the revision manuscript (cover letter). It is not necessary to accept all requests of the reviewers, but a clear explanation of why reviewers' comments were not accepted, has to be provided. If the deadline for re-submission is missed, the paper will be removed from the reviewing process.

The proof reading must be returned within three days. Only errors originating during preparation of the document for printing can be corrected. Standard proof marks will be used. No changes in the manuscript after acceptance for publication can be allowed.

Czech Journal of Animal Science **do not accept hard-copy paper manuscripts**; all manuscripts must be submitted electronically to the journal website. This method of submission results in much faster handling of your manuscript, fewer handling errors, and allows you to track the handling progress of your manuscript at any time. Manuscripts must be submitted as a Microsoft Word or other word processing document (filetype ".doc" or ".rtf").

Copyright. The journal is protected by copyright held by the publisher after the manuscript has been accepted for publication. As concerns the transfer of rights, the corresponding author takes over responsibility for all authors. No part of this publication may be reproduced, stored, or transmitted in any form or by any means, without the written permission of the publisher.

Manuscript layout. The Microsoft (MS) Word for Windows word-processing software should be used for creating the text in non-formatted style strictly following the journal layout. The manuscript must be typed (Times New Roman, font 12) with double spacing throughout. Lines must be numbered in a consecutive manner starting on the first page, in the left-hand margin. All pages of the manuscript must also be numbered consecutively.

If any abbreviations are used in the paper, they have to be explained appropriately when they are used in the text for the first time. It is not advisable to use any abbreviation in the paper title or in the abstract. Tables, graphs and other MS Word documents are to be submitted on separate pages appended to the article. The document must not be formatted in columns, heading styles, etc. This unique MS Word file must be saved under the first author's surname only. Graphs should be provided in MS Excel and they should be stored with original data. Photographs and autotypes should be submitted in high resolution (min. 300 dpi) JPG or TIFF format. All tables, graphs and photos should be numbered in the order in which they are included in the text, using Arabic numerals.

Manuscript length. Originals papers, including figures, tables and references, should not exceed 25 typed or computer-written pages. The number of figures and tables should be kept to a minimum.

The Title of the Paper should be short and informative, no subtitles or numbering the "serial" articles (Part I, Part II etc.) should be used.

The Abstract should not have more than 2,500 keystrokes (characters plus spaces). It should contain important information on methods used to solve the problem, clear description of results and their statistical significance, and brief and unambiguous conclusions drawn from the results. References and discussion of results should not be included in the abstract.

Keywords should not repeat nouns used in the title and should describe the studied problem as best as possible.

The Introduction section should provide information on the present state of research in the field concerned. It briefly justifies the research, specifies the hypotheses to be tested and gives the objective (s). References to literary sources document such

present findings that are used by the authors, not all that have been published until now. References in the text should agree with those in the list of references. It is recommended to include references to papers from peer periodicals only. Citations from non-available sources (reports, national journals, proceedings etc.) should be omitted. Papers published by one or two authors are to be cited by their names, those published by three or more authors by the name of the first one et al. If more than one paper by the same author(s) published in the same year and are cited, they should be differentiated by YEARa,b,c both in the text and the list of References. Names and year of publication are to be cited by including them in the text directly, e.g. "...as published by Brown (1995)" or indirectly – citing authors and year of publication in parenthesis (Green and Grey, 1996), (Jabl et al., 2002). Several papers cited together should be arranged according to the year of publication starting with the oldest one.

Material and Methods. All preliminary material, conducted experiments, their extent, conditions and course (experimental design) should be described in detail in this section. All original procedures that were used for the processing of experimental material and all analytical methods used for evaluation should also be detailed. Data verifying the quality of acquired data should be indicated for the used methods. The whole methodology is to be described only if it is an original one, in other cases it is sufficient to cite the author of the method and to mention any particular differences. Methods of statistical processing including the software used should also be listed in this section.

Results and Discussion. The results obtained from the experiments including their statistical evaluation and any commentary should be presented graphically or in tables in this section. The author should confront partial results with data published by other authors, whose names and year of publication are to be cited by including them in the text directly or indirectly.

References should be arranged in alphabetical order according to the surname and initials of authors. The year of publication cited in parenthesis, the full title of the paper in English with the language of publication in parenthesis, e.g. (in Czech) should follow. The title of the periodical should be typed in full.

Only papers cited in the text should be included in the list of references. All names of the authors must be printed in English transcription without national letters. Authors are responsible for the accuracy of their references.

Examples of references in the list:

- **Journal article**

Brown J. (1995): Estradiol determination in post-partum sows. *Journal of Endocrinology*, 198, 155–169.

- **Electronic journal article:**

Steinkraus K.H. (2002). Fermentation in world food processing. *Comprehensive Reviews in Food Science and Food Safety* [serial online]. 1, 23–32. Available from www.ift.org (accessed Apr 1, 2002).

- **Books and article within edited books, proceedings**

Spally M.R., Morgan S.S. (1989): *Methods of food analysis*. 2nd Ed. Elsevier, New York, 682 pp.

Kalab J. (1995): Changes in milk production during the sexual cycle. Hekel K. (ed.): *Lactation in Cattle*. Academic Press, London, UK, 876–888.

Janson L., Ahlin K.A. (1992): Postpartum reproductive performance in cattle selected for high and low fat content. In: *Proc. 43rd Annu. Mtg. European Association for Animal Production (EAAP)*. Madrid, Spain, 93–95.

- **Patent:**

Harred J.F., Knight A.R., McIntyre J.S., inventors; Dow Chemical Co., assignee. 1972 Apr 4. Epoxidation process. U.S. patent 3, 654, 317.

- **Dissertation:**

Smith D.E. (1988): Lipid oxidation at very low water activities. PhD Diss. Ithaca, NY: Cornell University. Available from: University Microfilms, Ann Arbor, MI: ABD62-83. 210 p.

Corresponding Author should include his or her full name including all academic, scientific and pedagogic titles and detailed address of the institution with postal code, phone and fax numbers, e-mail address. The author who is responsible for any correspondence with the journal should be indicated clearly.

Declaration of the authors must be carefully completed and signed by the first author, and then scanned in .pdf format to be uploaded in the submission of your manuscript, along side with the other manuscript files.

Offprints: Free reprint in Portable Document Format (PDF) sent via e-mail as an attachment.

Compliance with these instructions is obligatory for all authors. If a manuscript does not comply exactly with the above requirements, the editorial office will not accept it for a consideration and will return it to the authors without reviewing.

Revised February 2010