

## Effects of feeding corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> on performance and aflatoxin residues in broilers

J. YANG<sup>1</sup>, F. BAI<sup>1,2</sup>, K. ZHANG<sup>1</sup>, X. LV<sup>3</sup>, S. BAI<sup>1</sup>, L. ZHAO<sup>2</sup>, X. PENG<sup>1</sup>, X. DING<sup>1</sup>, Y. LI<sup>2</sup>, J. ZHANG<sup>2</sup>

<sup>1</sup>Key Laboratory of Animal Disease Resistance Nutrition of the Ministry of Education, Institute of Animal Nutrition, Sichuan Agricultural University, Sichuan Yaan, P.R.China

<sup>2</sup>Test Center for Feed Quality Supervision and Inspection (Chengdu) of the Ministry of Agriculture, Sichuan Chengdu, P.R.China

<sup>3</sup>Feed Safety Reference Laboratory of the Ministry of Agriculture, Feed Research Institute, Chinese Academy of Agricultural Sciences, Beijing, P.R.China

**ABSTRACT:** AFB<sub>1</sub> is the most abundant aflatoxin in food and animal feed, generally occurring along with low levels of other aflatoxins. In this experiment, broilers were administered corn that was naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>. We found that the broilers were more sensitive during the starter period and had a poor performance during the grower period. In addition, the broilers showed a significant reduction in feed intake when consuming material contaminated with 134.0 and 23.6 µg/kg of AFB<sub>1</sub> and AFB<sub>2</sub>, respectively. Our results also demonstrate that villus height and the ratio of villus height to crypt depth significantly decreased when the broilers were fed daily with diets contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>. Furthermore, AFB<sub>1</sub> residues in livers and breast muscles were determined by high-performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS) at levels of 0.137 and 0.016 µg/kg, respectively. The AFM<sub>1</sub> residue was also detected in livers at a level of 0.051 µg/kg but it was not detectable in breast muscles.

**Keywords:** mycotoxin; influence; poultry; HPLC-MS/MS

Mycotoxins are secondary toxic metabolites of filamentous fungi, which are harmful to human and animal health thus causing significant financial losses to animal industries (Hussein and Brasel, 2001; Wu and Munkvold, 2008; Zhang and Caupert, 2012). Aflatoxins (AF) naturally occur in moldy grains and are a group of metabolites produced by some strains of *Aspergillus flavus* and *Aspergillus parasiticus*. The primary aflatoxins of concern in feedstuffs are aflatoxins B<sub>1</sub> (AFB<sub>1</sub>), B<sub>2</sub> (AFB<sub>2</sub>), G<sub>1</sub> (AFG<sub>1</sub>), and G<sub>2</sub> (AFG<sub>2</sub>) (Monbaliu et al., 2010). The order of toxicity is AFB<sub>1</sub> > AFG<sub>1</sub> > AFB<sub>2</sub> > AFG<sub>2</sub>.

The legal limits of AFB<sub>1</sub> and AFT in total for several agricultural products have been established by the European Union (2–12 µg/kg for AFB<sub>1</sub>, 4–15 µg/kg for aflatoxins total (B<sub>1</sub> + G<sub>1</sub> + B<sub>2</sub> + G<sub>2</sub>), and 0.05 µg/kg for AFM<sub>1</sub> in milk), but the value of aflatoxins in other animal products was not recommended (European Commission, 2010).

AFB<sub>1</sub> is transformed into toxic metabolites primarily in the liver, and the metabolite of AFB<sub>1</sub>-8, 9-epoxide can form covalent bonds with DNA, RNA, and proteins, resulting in a potent carcinogenic effect (Swenson et al., 1977; Nomura et al.,

Supported by the specific research supporting program for academic sustenance research team in the Sichuan Agricultural University and the specific research and the Feed Safety project of the Agricultural Ministry of China.

2011). AFB<sub>2</sub> is converted to AFB<sub>1</sub> in the livers of both rats and ducks, and the rates of these reactions appear to correlate with the toxic potential of AFB<sub>2</sub> in these species (Wogan et al., 1971). AFM<sub>1</sub> is the toxic metabolite of AFB<sub>1</sub>, and AFM<sub>2</sub> is the hydroxylated form of AFB<sub>2</sub> (Wolzak et al., 1985; Bintvihok et al., 2003; Hussain et al., 2010; Bianco et al., 2012). AFM<sub>1</sub> and AFM<sub>2</sub> are commonly associated with milk and other edible animal products.

AFB<sub>1</sub> contaminants occur more frequently at higher concentrations than the other AF metabolites (Fandohan et al., 2005), and AFB<sub>1</sub> toxicity in poultry has been well elucidated (Martinez-de-Anda et al., 2010). Previous studies have shown that AFB<sub>1</sub> has a range of negative effects on poultry health, including poor performance (Oguz and Kurtoglu, 2000; Magnoli et al., 2011), liver and kidney lesions (Ortatatli et al., 2005), immunity damage (Marin et al., 2002), and hematology problems (Zhao et al., 2010). Moreover, the tissue deposition and residue of AFB<sub>1</sub> in poultry has also been determined (Oliveira et al., 2000, 2003). However, AFB<sub>1</sub> usually occurs in feedstuff along with low levels of other aflatoxins, and the toxicity observed in poultry is complex and diverse (Ghahri et al., 2010; Yunus et al., 2011).

AFB<sub>1</sub> is commonly found along with AFB<sub>2</sub> in corn in the southwest of China (Placinta et al., 1999), but few studies have explored the performance of broilers and the level of aflatoxin residues in the tissues of broilers administered with naturally occurring AFB<sub>1</sub> and AFB<sub>2</sub> grains. In this study, we explored the growth performance and nutrient digestibility of broilers and determined the aflatoxin residue levels in broilers fed with corn that was naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>.

## MATERIAL AND METHODS

### Animals and management

A total of 1200 one-day-old COBB500 male broilers were weighed and randomly allocated into 5 treatment groups. Each treatment group had 8 replicates of 30 broilers. All the ingredients in the 5 treatment diets, except for the corn, were the same. The basal control diet was formulated and compounded to meet the nutritional requirements of commercial broilers (National Research Council, 1994) during the starter and grower periods, presented in Table 1. The basal control diet

was not contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>. The control group (I) was fed the basal control diet. The second (II), third (III), fourth (IV), and fifth groups (V) were given feed in which the ratio of naturally contaminated corn as a substitute for normal corn was 25, 50, 75, and 100%, respectively. The study

Table 1. Composition and nutrient levels of the basal diets (%)

Items	Day 1–21	Day 22–42
<b>Ingredient</b>		
Corn	60.00	63.00
Soybean meal (46%)	31.20	28.17
Corn gluten meal	3.00	2.50
Soybean oil	1.14	2.27
Calcium carbonate	0.95	0.88
Calcium hydrophosphate	1.85	1.60
L-Lysine hydrochloride	0.08	0.02
DL-Methionine	0.16	0.09
Salt	0.50	0.40
Choline chloride	0.15	0.10
Vitamin and mineral premix	0.33 <sup>a</sup>	0.32 <sup>b</sup>
Rice hull	0.64	0.65
Total	100.00	100.00
<b>Nutrient levels</b>		
Crude protein	20.83	19.37
ME (MJ/kg)	12.15	12.55
Calcium	0.97	0.87
Available phosphorus	0.44	0.39
Lysine	1.11	0.97
Methionine	0.48	0.39
Methionine and cysteine	0.86	0.72
Threonine	0.78	0.74

<sup>a</sup>provided per kg of diet: vitamin A 12 000 IU, cholecalciferol 3000 IU, vitamin E 7.5 IU, vitamin K<sub>3</sub> 1.5 mg, thiamin 0.6 mg, riboflavin 4.8 mg, pyridoxine 1.8 mg, vitamin B12 9 µg, folic acid 150 µg, niacin 10.5 mg, calcium pantothenic acid 7.5 mg, iron 100 mg, copper 8 mg, manganese 120 mg, zinc 100 mg, selenium 0.3 mg, iodine 0.7 mg

<sup>b</sup>provided per kg of diet: vitamin A 8000 IU, cholecalciferol 2000 IU, vitamin E 5 IU, vitamin K<sub>3</sub> 1 mg, thiamin 0.4 mg, riboflavin 3.2 mg, pyridoxine 1.2 mg, vitamin B<sub>12</sub> 6 µg, folic acid 100 µg, niacin 7 mg, calcium pantothenic acid 5 mg, iron 80 mg, copper 8 mg, manganese 100 mg, zinc 80 mg, selenium 0.3 mg, iodine 0.7 mg

protocol was conducted in accordance with the animal husbandry laws of the P.R.China.

The broilers were fed in cage pens for 42 days and maintained on a 24-h continuous light schedule. The initial temperature of  $33 \pm 1^\circ\text{C}$  was gradually reduced by  $2^\circ\text{C}$  per week to reach  $21^\circ\text{C}$  after 35 days, and this temperature was maintained for the rest of the experiment. Feed and water were provided *ad libitum*.

### Analysis of dietary mycotoxins

The corn was ground in a Restch ZM100 mill using a 0.75 mm sieve. Mycotoxins were extracted from 25 g of a milled sample with 50 ml of water for deoxynivalenol (DON), acetonitrile and water (90 : 10, v/v) solution for zearalenone (ZEA), methanol and water (70 : 30, v/v) solution for AF, and methanol and water (80 : 20, v/v) solution for T-2 toxin, ochratoxin A (OTA), and fumonisin B<sub>1</sub> (FB<sub>1</sub>). The aforementioned solutions were shaken vigorously in a sealed flask for 40 min. The extract was filtered by gravity, and then 10 ml of the filtrate was mixed with 40 ml of PBS. The mixture was centrifuged at 8000 g for 15 min. Then, 15 ml of the supernatant was subjected to a purification process, which was carried out with an AflaStar<sup>TM</sup> Immunoaffinity columns (Romer Labs Diagnostic GmbH, Tulln, Austria). After the sample had passed through the column, the column was washed with 10 ml of PBS and 10 ml of water and then it was eluted with 2 ml of methanol into a glass vial, which was maintained at  $4^\circ\text{C}$  until analysis.

The mycotoxins were quantified via high-performance liquid chromatography equipped with a fluorescence detector Agilent 1100 (Agilent Technologies Inc., Santa Clara, USA). Separation was achieved on a SB-C<sub>18</sub> column (4.6 × 250 mm, 5 μm) (Agilent Technologies Inc., Santa Clara, USA). The mobile phases were mixtures of methanol and water (45 : 55, v/v) solution for AF, acetonitrile and water (12 : 88, v/v) solution for DON, acetonitrile and water (80 : 20, v/v) solution for T-2 toxin, acetonitrile and water and methanol (46 : 46 : 8, v/v/v) solution for ZEA, methanol and aqueous 1% acetic acid (50 : 50, v/v) solution for OTA, and methanol and water acidified with 0.1 mol/l phosphoric acid (pH = 3.3) (50 : 50, v/v) solution for OTA. The flow rate was 1 ml/min, and the injection volume was 50 μl.

A post-column photochemical derivatization was used to enhance the mycotoxin response using a

PHRED photochemical reactor (AURA Industries Inc., New York, USA). The wavelengths of excitation and emission were fixed at 365 and 440 nm for AF, 330 and 460 nm for OTA, 330 and 440 nm for FB<sub>1</sub>, and 381 and 470 nm for T-2 toxin, respectively. However, the presence of ZEA and DON was monitored at 274 and 220 nm with the fluorescence detector, respectively. The above mycotoxins were analyzed by the Test Center for Feed Quality Supervision and Inspection (Chengdu) of the Ministry of Agriculture, with detection limits (in μg/kg) of 2 for AFB<sub>1</sub>, 0.8 for AFB<sub>2</sub>, 2.5 for AFG<sub>1</sub>, 1.5 for AFG<sub>2</sub>, 100 for T-2 toxin, 300 for DON, 100 for ZEN, 30 for OTA, and 200 for FB<sub>1</sub> (Administration of Quality and Technology Supervision of Sichuan Province, 2010).

### Body weight and feed consumption

Broilers were weighed individually, and feed consumption for each pen was measured on days 22 and 43 during the 6-week experiment. The cumulative weight gain and feed consumption were determined, and the weekly and cumulative gain : feed ratios were calculated.

### Digestibility experiment

The excreta collection method was used to explore nutrient digestibility. The experiment consisted of a 2-day adaptation period and 4-day collection period. At day 43, one broiler per pen was randomly separated from the group and kept in isolation. Feed and water were provided *ad libitum*. Excreta were collected every 6 h and stored at  $-20^\circ\text{C}$ . At the end of the collection period, the feed was collected and weighed.

Feed and excreta samples were subsequently analyzed for dry matter, organic matter, crude protein, calcium, and phosphorus according to the AOAC Standard Methods (AOAC, 1980). Moreover, the gross energy was determined by adiabatic bomb calorimetry according to instructions provided by the manufacturer (Parr Instrument Company, Moline, USA).

### Intestinal morphology

On days 22 and 43, histomorphological samples were taken from the duodenum, jejunum, and

ileum. The samples were fixed in 10% formalin. Then, the fixed tissues were washed, dehydrated, trimmed, and embedded in paraffin. Sections of 5  $\mu\text{m}$  were sliced, mounted on slides, and then stained with hematoxylin and eosin for histomorphological analysis. For each section, after taking a picture ( $4 \times 10$ ) with an Olympus CX31-32C02 optical microscope (Olympus Optical Co., Ltd., Tokyo, Japan), Image-Pro-Plus 6.0 was used to measure 8 of the typical villi from crypt mouth to villus tip and 8 of the associated crypts from crypt mouth to base. The ratio of villus height to crypt depth was calculated.

### Sample preparation for HPLC-MS/MS

Eight samples of liver and breast muscle were collected from groups I, III, and V. The samples were homogenized with IKA ULTRA-TURRAX T 10 disperser (Sigma-Aldrich Corporation, St. Louis, USA). 10 g of the tissue sample was weighed and mixed with 1 ml of 20% aqueous citric acid and 2 g diatomaceous earth. The mixture was sonicated with 20 ml of dichloromethane for 40 min. After centrifugation at 11 000  $g$  for 10 min, 10 ml of the supernatant was evaporated in a vacuum (Tavcar-Kalcher et al., 2007). The dry medium residue was dissolved with 3 ml of methanol and mixed with 9 ml of PBS solution. A total of 4 ml of the solution was applied to the AflaStar<sup>TM</sup> R Immunoaffinity columns (Romer Labs Diagnostic GmbH, Tulln, Austria), eluted with 3 ml of methanol, and dried by nitrogen. Then, 0.5 ml of methanol and 0.1% formic acid (30 : 70, v/v) was added and filtered through a 0.22- $\mu\text{m}$  filter into a 2-ml autosampler vial.

### Quantification of AFB<sub>1</sub> and AFM<sub>1</sub> residues in tissues with HPLC-MS/MS

An Agilent 1290 infinity series HPLC system with an SB-C18 column ( $2.1 \times 50$  mm) (Agilent Technologies Inc., Santa Clara, USA) was used. The column oven was set at 30°C. The gradient elution used two solutions: solution A was methanol and solution B was 0.1% formic acid in water. The gradient elution program was as follows: solution A was changed from 30% at 0 min to 90% at 5 min and from 90% at 5 min to 30% at 9 min. A flow rate of 0.3 ml/min was used.

Mass spectral analysis was performed using an Agilent 6460 Triple Quad LC/MS (Agilent Technologies Inc., Santa Clara, USA) fitted to an electrospray probe operating in the positive mode (ESI<sup>+</sup>). The potential set on the capillary was 4 kV. The nitrogen-gas flow rates for nebulization and desolvation were 10 l/min and 35 psi, respectively. The pressure of the collision gas of nitrogen was 0.2 MPa. The signal acquisition was performed in Multiple Reaction Monitoring Mode (MRM). Precursor ions for AFB<sub>1</sub> and AFM<sub>1</sub> were  $m/z = 313$  and 329, respectively. The product ions for the target compounds were  $m/z = 269, 241$  for AFB<sub>1</sub> and 273, 259 for AFM<sub>1</sub>. The quantitation ion transitions were  $m/z = 313, 241$  for AFB<sub>1</sub> and 329, 273 for AFM<sub>1</sub>.

Standard solutions of AFB<sub>1</sub> and AFM<sub>1</sub> were purchased from Sigma-Aldrich Corporation (St. Louis, USA). Because single point calibration was used, three concentrations of the standard solution of AFB<sub>1</sub> and AFM<sub>1</sub> in the liver were prepared at 0.04, 0.08, and 0.16  $\mu\text{g/l}$  by diluting with methanol and 0.1% formic acid (30 : 70, v/v). Moreover, blank broiler tissues used for method validation were first analyzed, and no aflatoxin residue was detected. Then, 10-g portions of tissue homogenate were fortified to produce samples ( $n = 5$ ) containing 0.01, 0.02, and 0.05  $\mu\text{g/kg}$  of AFB<sub>1</sub> and AFM<sub>1</sub>. The limit of detection (LOD) and limit of quantification (LOQ) were considered to be the concentrations in tissue samples that produced signal-to-noise (S/N) ratios of 3 and 10, respectively.

### Statistical analysis

Data were analyzed by One-Way ANOVA using SPSS software, Version 11.0. When the ANOVA showed significance, Duncan's significant-difference test was applied. Data of AFB<sub>1</sub> and AFM<sub>1</sub> residues were analyzed using Independent Samples  $t$ -Tests. All statements of differences were based on a significance of  $P < 0.05$ .

## RESULTS AND DISCUSSION

### Dietary mycotoxin concentrations

The corn used in this study was mainly contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>. The AFB<sub>1</sub> contents in diets were 16.3–82.4  $\mu\text{g/kg}$  in the starter period and

Table 2. Concentrations of mycotoxins in diet and corn ( $\mu\text{g/kg}$ )

Diet	I	II	III	IV	V	Control corn	Contaminated corn
<b>Day 1–21</b>							
AFB <sub>1</sub>	nd	16.3	36.9	45.6	82.4	nd	149.6
AFB <sub>2</sub>	nd	3.2	6.4	7.9	14.2	nd	24.2
<b>Day 22–42</b>							
AFB <sub>1</sub>	nd	34.3	69.3	95.2	134.0	nd	229.0
AFB <sub>2</sub>	nd	6.2	12.1	17.0	23.6	nd	37.8

nd = not detectable

34.3–134  $\mu\text{g/kg}$  in the grower period (Table 2). The AFB<sub>2</sub> concentrations in diets were 3.2–14.2 and 6.2–23.6  $\mu\text{g/kg}$  in the starter and grower periods, respectively. The contents of other mycotoxins, including AFG<sub>1</sub>, AFG<sub>2</sub>, DON, ZEA, OTA, T-2 toxin, and FB<sub>1</sub> were below the limit of detection.

### Body weight and feed consumption

Compared with control, the body weight gain and feed consumption of broilers, which were fed diets contaminated with more than 36.9 and 6.4  $\mu\text{g/kg}$  of AFB<sub>1</sub> and AFB<sub>2</sub>, significantly decreased ( $P < 0.05$ )

from days 1 to 21. The feed : gain ratio significantly increased ( $P < 0.05$ ) when the broilers were fed diets contaminated with 82.4 and 14.2  $\mu\text{g/kg}$  of AFB<sub>1</sub> and AFB<sub>2</sub> (Table 3). During days 22 to 42, only broilers that consumed diets contaminated with 134 and 23.6  $\mu\text{g/kg}$  of AFB<sub>1</sub> and AFB<sub>2</sub> showed a significant reduction in feed intake. We also found that feed consumption significantly decreased over the full period for groups IV and V ( $P < 0.05$ ), and body weight gain significantly decreased over the full period for group V ( $P < 0.05$ ).

When the broilers were administered feed contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>, our results prove that some economically significant effects occurred

Table 3. Effect of corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> on performance of broilers

Diet	I	II	III	IV	V	<i>P</i>
<b>Body weight (BW)</b>						
Day 21	828.7 $\pm$ 7.5 <sup>a</sup>	831.1 $\pm$ 22.1 <sup>a</sup>	807.8 $\pm$ 12.0 <sup>b</sup>	788.7 $\pm$ 15.4 <sup>c</sup>	749.4 $\pm$ 17.4 <sup>d</sup>	< 0.001
Day 42	2434.8 $\pm$ 120.0 <sup>ab</sup>	2417.9 $\pm$ 113.8 <sup>ab</sup>	2469.7 $\pm$ 102.7 <sup>a</sup>	2349.6 $\pm$ 75.0 <sup>bc</sup>	2301.3 $\pm$ 69.3 <sup>c</sup>	0.011
<b>Body weight gain (BWG)</b>						
Day 1–21	788.2 $\pm$ 7.5 <sup>a</sup>	790.6 $\pm$ 22.1 <sup>a</sup>	767.3 $\pm$ 12.0 <sup>b</sup>	748.2 $\pm$ 15.4 <sup>c</sup>	708.9 $\pm$ 17.4 <sup>d</sup>	< 0.001
Day 22–42	1606.0 $\pm$ 115.8	1586.8 $\pm$ 110.8	1661.9 $\pm$ 105.5	1561.0 $\pm$ 73.5	1551.9 $\pm$ 58.8	0.177
Day 1–42	2394.2 $\pm$ 120.2 <sup>ab</sup>	2377.4 $\pm$ 113.8 <sup>ab</sup>	2429.2 $\pm$ 102.7 <sup>a</sup>	2309.1 $\pm$ 75.0 <sup>bc</sup>	2260.8 $\pm$ 69.3 <sup>c</sup>	0.011
<b>Feed consumption (FC)</b>						
Day 1–21	1134.7 $\pm$ 13.8 <sup>a</sup>	1134.4 $\pm$ 24.7 <sup>a</sup>	1109.1 $\pm$ 11.7 <sup>b</sup>	1091.0 $\pm$ 14.0 <sup>b</sup>	1050.7 $\pm$ 31.8 <sup>c</sup>	< 0.001
Day 22–42	3414.2 $\pm$ 205.9 <sup>ab</sup>	3345.0 $\pm$ 223.7 <sup>abc</sup>	3456.0 $\pm$ 98.3 <sup>a</sup>	3250.4 $\pm$ 176.8 <sup>bc</sup>	3221.1 $\pm$ 129.5 <sup>c</sup>	0.044
Day 1–42	4377.7 $\pm$ 201.0 <sup>a</sup>	4312.1 $\pm$ 210.1 <sup>ab</sup>	4416.7 $\pm$ 108.1 <sup>a</sup>	4179.0 $\pm$ 151.5 <sup>bc</sup>	4131.6 $\pm$ 148.6 <sup>c</sup>	0.006
<b>FC/BWG</b>						
Day 1–21	1.44 $\pm$ 0.02 <sup>bc</sup>	1.44 $\pm$ 0.02 <sup>c</sup>	1.45 $\pm$ 0.02 <sup>bc</sup>	1.46 $\pm$ 0.02 <sup>b</sup>	1.48 $\pm$ 0.01 <sup>a</sup>	< 0.001
Day 22–42	2.13 $\pm$ 0.08	2.11 $\pm$ 0.07	2.09 $\pm$ 0.09	2.08 $\pm$ 0.09	2.07 $\pm$ 0.06	0.647
Day 1–42	1.83 $\pm$ 0.04	1.82 $\pm$ 0.03	1.82 $\pm$ 0.05	1.81 $\pm$ 0.05	1.83 $\pm$ 0.03	0.891

<sup>a–d</sup>different lowercases in the same row mean significant difference between treatments ( $P < 0.05$ )

data are mean (g)  $\pm$  SD ( $n = 8$ )

in the broilers, including reduced body weight gain, feed consumption, and feed conversion rate. Aravind et al. (2003) reported that grains naturally contaminated with mycotoxins (AF 168 µg/kg, OAT 8.4 µg/kg, ZEN 54 µg/kg, T-2 toxin 32 µg/kg) significantly resulted in poor performance. Although Magnoli et al. (2011) found that diets containing 50 µg/kg AFB<sub>1</sub> had no effect on the performance of broilers from 28 to 46 days of age, our results prove that broilers were more sensitive to aflatoxins during the starter period, while broilers administered AFB<sub>1</sub>- and AFB<sub>2</sub>-contaminated diets showed a poor performance during the grower period.

### Nutrient digestibility

Our results prove that corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> had no effect on the apparent digestibility of calcium and phosphorus ( $P > 0.05$ ). Our results are in agreement with the results of Kermanshahi et al. (2007) in broilers and Applegate et al. (2009) in laying hens. However, there was a trend that showed a decrease in the apparent digestibility of dry matter, organic matter, crude protein, and gross energy of the broilers feed diets when contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> ( $P > 0.05$ ). We found that dry matter, organic matter, crude protein, and gross energy decreased from 72.5, 75.1, 51.5, and 75.6% to 69.4, 71.8, 42.9, and

72.8%, respectively. Han et al. (2008) have reported that the 20- and 40-µg/kg AFB<sub>1</sub> diets reduced the apparent digestibility of crude protein but had no effect on the apparent digestibility of ether extract, ash, calcium or total phosphorus.

### Intestinal morphology

Tables 4 and 5 show the results of intestinal morphology. During the period from day 1 to day 21 of age, corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> significantly decreased the villus height in the duodenum ( $P < 0.05$ ). The ratio of villus height to crypt depth in the duodenum decreased with increasing content of contaminated corn in the diets, and the same trend was observed in the jejunum and ileum. During the period from day 22 to day 42 of age, the villus height in the duodenum decreased with increasing content of contaminated corn in the diets, and the trend for crypt depth was just the opposite. Moreover, we found that the ratio of villus height to crypt depth in the duodenum showed a significant decrease ( $P < 0.05$ ) when the broilers consumed a diet contaminated with more than 95.2 and 17.0 µg/kg of AFB<sub>1</sub> and AFB<sub>2</sub>, respectively. The same trend was found in the jejunum and ileum results.

Increasing of the villus height would suggest an increased surface area capable of greater absorption of available nutrients. The ratio of villus

Table 4. Effect of corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> on intestinal morphology of broilers on day 21

Diet	I	II	III	IV	V	P
<b>Duodenum (µm)</b>						
Villus height	1479.5 ± 140.4 <sup>a</sup>	1202.4 ± 54.3 <sup>b</sup>	1263.4 ± 151.7 <sup>b</sup>	1210.9 ± 134.3 <sup>b</sup>	996.7 ± 145.7 <sup>c</sup>	< 0.001
Crypt depth	214.6 ± 23.8	182.4 ± 44.9	219.4 ± 36.7	207.7 ± 16.9	211.9 ± 12.8	0.32
V/C	6.96 ± 0.97 <sup>a</sup>	6.98 ± 2.06 <sup>a</sup>	5.86 ± 1.05 <sup>ab</sup>	5.85 ± 0.65 <sup>ab</sup>	4.73 ± 0.89 <sup>b</sup>	0.037
<b>Jejunum (µm)</b>						
Villus height	1172.3 ± 150.6	1057.3 ± 222.8	961.1 ± 70.4	1033.0 ± 146.6	989.6 ± 203.1	0.273
Crypt depth	213.4 ± 54.7	251.2 ± 43.0	288.0 ± 13.7	257.8 ± 56.9	242.2 ± 40.0	0.091
V/C	5.74 ± 1.37 <sup>a</sup>	4.32 ± 1.23 <sup>b</sup>	3.35 ± 0.36 <sup>b</sup>	4.12 ± 0.79 <sup>b</sup>	4.14 ± 0.90 <sup>b</sup>	0.008
<b>Ileum (µm)</b>						
Villus height	877.8 ± 112.1	895.3 ± 139.1	825.9 ± 99.7	793.0 ± 36.9	749.9 ± 21.5	0.069
Crypt depth	259.6 ± 34.7	222.3 ± 30.2	248.2 ± 48.5	281.3 ± 74.0	296.3 ± 57.6	0.115
V/C	3.43 ± 0.62 <sup>ab</sup>	4.09 ± 0.80 <sup>a</sup>	3.45 ± 0.91 <sup>ab</sup>	2.98 ± 0.72 <sup>b</sup>	2.61 ± 0.45 <sup>b</sup>	0.013

<sup>a-c</sup>different lowercases in the same row mean significant difference between treatments ( $P < 0.05$ )

V/C = villus height/crypt depth

data are mean ± SD ( $n = 8$ )

Table 5. Effect of corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> on intestinal morphology of broilers on day 42

Diet	I	II	III	IV	V	P
<b>Duodenum (μm)</b>						
Villus height	1445.5 ± 175.9 <sup>ab</sup>	1481.1 ± 111.4 <sup>a</sup>	1540.7 ± 123.1 <sup>a</sup>	1359.1 ± 190.7 <sup>ab</sup>	1270.9 ± 89.0 <sup>b</sup>	0.047
Crypt depth	182.4 ± 20.7 <sup>b</sup>	237.2 ± 34.3 <sup>a</sup>	225.1 ± 22.9 <sup>a</sup>	259.9 ± 48.2 <sup>a</sup>	230.6 ± 23.3 <sup>a</sup>	0.007
V/C	7.45 ± 1.62 <sup>a</sup>	6.40 ± 1.30 <sup>ab</sup>	6.86 ± 0.22 <sup>ab</sup>	5.35 ± 1.01 <sup>b</sup>	5.58 ± 0.91 <sup>b</sup>	0.029
<b>Jejunum (μm)</b>						
Villus height	1238.0 ± 204.4	995.7 ± 171.1	1119.3 ± 206.1	1083.8 ± 199.6	1002.2 ± 129.1	0.190
Crypt depth	181.4 ± 26.3	185.8 ± 46.5	211.9 ± 51.0	219.0 ± 55.2	216.4 ± 43.2	0.487
V/C	6.87 ± 1.04 <sup>a</sup>	5.53 ± 1.25 <sup>b</sup>	5.41 ± 0.98 <sup>b</sup>	4.64 ± 0.67 <sup>b</sup>	4.77 ± 1.11 <sup>b</sup>	0.007
<b>Ileum (μm)</b>						
Villus height	785.5 ± 104.7	780.3 ± 109.1	800.3 ± 130.6	701.8 ± 236.4	609.6 ± 53.0	0.158
Crypt depth	136.7 ± 20.0	156.7 ± 29.9	181.9 ± 43.9	176.1 ± 56.3	142.1 ± 17.4	0.200
V/C	5.67 ± 0.86 <sup>a</sup>	5.22 ± 1.09 <sup>ab</sup>	4.56 ± 1.11 <sup>bc</sup>	4.02 ± 0.52 <sup>c</sup>	4.32 ± 0.46 <sup>bc</sup>	0.016

<sup>a-c</sup>different lowercases in the same row mean significant difference between treatments ( $P < 0.05$ )

V/C = villus height/crypt depth

data are mean ± SD ( $n = 8$ )

height to crypt depth might reflect differences in the digestion and absorption of the small intestines (Caspary, 1992). A previous study has demonstrated that grains naturally contaminated with DON significantly reduced the height, width, and surface of villus in the duodenum and jejunum of broilers (Girish and Smith, 2008). However, just few reports in the literature regarding the effects of AFB<sub>1</sub> and AFB<sub>2</sub> on the intestinal morphology have been published. Our results demonstrate that the villus height and the ratio of villus height to crypt depth significantly decreased when the broilers were fed daily with diets contaminated with AFB<sub>1</sub> and AFB<sub>2</sub>. We also found that the mixture of AFB<sub>1</sub> and AFB<sub>2</sub> mainly affected the morphology of the duodenum, indicating that long-term exposure to AFB<sub>1</sub> and AFB<sub>2</sub> would stimulate the proximal gastrointestinal tract, affect the characteristics of the intestine morphology, and alter the absorption of nutrients.

### Method performance for HPLC-MS/MS

The techniques currently used to detect and quantify AFB<sub>1</sub> and AFM<sub>1</sub> include ELISA (Reddy et al., 2009; Yang et al., 2009), HPLC (Chen et al., 2005; Meucci et al., 2010), and HPLC-MS/MS (Ediage et al., 2011; Van Pamel et al., 2011). The HPLC-MS/MS-based method was selected in this work because AFB<sub>1</sub> and AFM<sub>1</sub> were detected in trace analyses. The peak profile of the AFB<sub>1</sub> and AFM<sub>1</sub>

standard was similar to the profile of AFB<sub>1</sub> and AFM<sub>1</sub> from the liver sample (data not shown).

In our studies, the recoveries of AFB<sub>1</sub> and AFM<sub>1</sub> from fortified samples were 70.8–85.7% and 68.9–83.0% over the concentration range of 0.01 to 0.05 μg/kg. For each calibration point, the values of the relative standard deviation (RSD) ranged from 1.9 to 10.5%. The LOD for AFB<sub>1</sub> and AFM<sub>1</sub> in tissue samples, defined as the concentrations that produced the S/N ratio of 3, was 0.005 μg/kg. The LOQ for AFB<sub>1</sub> and AFM<sub>1</sub>, defined as the concentration that produced the S/N ratio of 10, was 0.02 μg/kg in tissue samples. The quantification and detection limits (0.02 and 0.005 μg/kg) of our analytical technique were similar to the results of Magnoli et al. (2011), who detected AFB<sub>1</sub> residues in livers by the HPLC-MS/MS method.

### AFB<sub>1</sub> and AFM<sub>1</sub> residues in tissues

The levels of AFB<sub>1</sub> and AFM<sub>1</sub> in the livers from group V were significantly higher than those of the livers from group III (Table 6). The levels of AFB<sub>1</sub> residue in the livers from groups III and V (0.049 and 0.137 μg/kg, respectively) were higher than the levels in breast muscles. In a previous study, high levels of AFB<sub>1</sub> residues in livers (3.51–6.97 μg/kg) and breast muscles (1.63–3.27 μg/kg) were detected when broilers were given feed containing 1.6–6.4 mg/kg AFB<sub>1</sub> (Hussain et al., 2010). Residue

Table 6. Concentrations of AFB<sub>1</sub> and AFM<sub>1</sub> residues in tissues of broilers (µg/kg)

Diet		Control	III	V	P
AFB <sub>1</sub>	liver	nd	0.049 ± 0.025 <sup>a</sup>	0.137 ± 0.037 <sup>b</sup>	0.001
	muscle	nd	0.015 ± 0.003	0.016 ± 0.007	0.671
AFM <sub>1</sub>	liver	nd	0.033 ± 0.006 <sup>a</sup>	0.051 ± 0.009 <sup>b</sup>	0.004
	muscle	nd	nd	nd	

<sup>a,b</sup>different lowercases in the same row mean significant difference between treatments ( $P < 0.05$ )

nd = not detectable

data are mean ± SD ( $n = 8$ )

levels of AFB<sub>1</sub> and AFM<sub>1</sub> were also detected in the kidneys and thighs of both male broilers and hens that were fed a diet contaminated with 50 µg/kg of AFB<sub>1</sub> (Micco et al., 1988). Moreover, Bintvihok and Kositcharoenkul (2006) found AFB<sub>1</sub> residues at levels of 0.13 and 0.02 µg/kg and the AFM<sub>1</sub> residues at levels of 0.32 and 0.08 µg/kg, respectively in the livers and muscles of broilers receiving feed contaminated with 100 µg/kg of AFB<sub>1</sub>. Meanwhile, in hens that were administered a diet containing 3 mg/kg AFB<sub>1</sub>, the ratios of AFB<sub>1</sub> in the feed to the residual level in the egg yolk and albumen were 4615 and 3846, respectively (Bintvihok et al., 2002). In our study, the concentrations of AFM<sub>1</sub> residue in the livers from groups III and V were 0.033 and 0.051 µg/kg, respectively. However, the levels of AFM<sub>1</sub> residue in breast muscle were not detectable.

In conclusion, our results indicate that corn naturally contaminated with AFB<sub>1</sub> and AFB<sub>2</sub> decreased growth performance and altered intestine morphology in broilers. The residues of AFB<sub>1</sub> and AFM<sub>1</sub> were detected in the edible tissues of broilers, although the animal feedstuff contained low levels of AFB<sub>1</sub> and AFB<sub>2</sub>.

## Acknowledgement

We are grateful to the financial support of the specific research program for academic sustenance of research teams at Sichuan Agricultural University and the safety of mycotoxin warning program of the Agricultural Ministry of China.

## REFERENCES

AOAC (1980): Official Methods of Analysis. 13<sup>th</sup> Ed. Association of Official Analytical Chemists, Washington, USA.

Applegate T., Schatzmayr G., Prickett K., Troche C., Jiang Z. (2009): Effect of aflatoxin culture on intestinal function and nutrient loss in laying hens. *Poultry Science*, 88, 1235–1241.

Administration of Quality and Technology Supervision of Sichuan Province (2010): Sichuan Provincial Standard for the determination of aflatoxins B1, B2, G1, G2 (DB 51/T 1077-2010), T-2 toxin (DB 51/T 1078-2010), deoxynivalenol (DB 51/T 1079-2010), ochratoxin (DB 51/T 1081-2010), fumonisin B1 (DB 51/T 1082-2010) in feeds with HPLC.

Aravind K.L., Patil V.S., Devegowda G., Umakantha B., Ganpule S.P. (2003): Efficacy of esterified glucomannan to counteract mycotoxicosis in naturally contaminated feed on performance and serum biochemical and hematological parameters in broilers. *Poultry Science*, 82, 571–576.

Bianco G., Russo R., Marzocco S., Velotto S., Autore G., Severino L. (2012): Modulation of macrophage activity by aflatoxins B1 and B2 and their metabolites aflatoxins M1 and M2. *Toxicon*, 59, 644–650.

Bintvihok A., Kositcharoenkul S. (2006): Effect of dietary calcium propionate on performance, hepatic enzyme activities and aflatoxin residues in broilers fed a diet containing low levels of aflatoxin B1. *Toxicon*, 47, 41–46.

Bintvihok A., Ponpornpisit A., Tangtrongpiros J., Panichkriangkrai W., Rattanapanee R., Doi K., Kumagai S. (2003): Aflatoxin contamination in shrimp feed and effects of aflatoxin addition to feed on shrimp production. *Journal of Food Protection*, 66, 882–885.

Bintvihok A., Thiengnin S., Doi K., Kumagai S. (2002): Residues of aflatoxins in the liver, muscle and eggs of domestic fowls. *The Journal of Veterinary Medical Science*, 64, 1037–1039.

Caspary W.F. (1992): Physiology and pathophysiology of intestinal absorption. *The American Journal of Clinical Nutrition*, 55, 299S–308S.

Chen C.Y., Li W.J., Peng K.Y. (2005): Determination of aflatoxin M1 in milk and milk powder using high-flow solid-phase extraction and liquid chromatography-



- tandem mass spectrometry. *Journal of Agricultural and Food Chemistry*, 53, 8474–8480.
- Ediage E.N., Di Mavungu J.D., Monbaliu S., Van Peteghem C., De Saeger S. (2011): A validated multianalyte LC-MS/MS method for quantification of 25 mycotoxins in cassava flour, peanut cake and maize samples. *Journal of Agricultural and Food Chemistry*, 59, 5173–5180.
- European Commission (2010): Commission Regulation (EU) No. 165/2010 of 26 February 2010 amending Regulation (EC) No. 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards aflatoxins. *Official Journal of the European Union*, L50, 8–12.
- Fandohan P., Zoumenou D., Hounhouigan D.J., Marasas W.F., Wingfield M.J., Hell K. (2005): Fate of aflatoxins and fumonisins during the processing of maize into food products in Benin. *International Journal of Food Microbiology*, 98, 249–259.
- Ghahri H., Habibian R., Fam M.A. (2010): Evaluation of the efficacy of esterified glucomannan, sodium bentonite, and humic acid to ameliorate the toxic effects of aflatoxin in broilers. *Turkish Journal of Veterinary and Animal Sciences*, 34, 385–391.
- Girish C., Smith T. (2008): Effects of feeding blends of grains naturally contaminated with fusarium mycotoxins on small intestinal morphology of turkeys. *Poultry Science*, 87, 1075–1082.
- Han X.Y., Huang Q.C., Li W.F., Jiang J.F., Xu Z.R. (2008): Changes in growth performance, digestive enzyme activities and nutrient digestibility of cherry valley ducks in response to aflatoxin B1 levels. *Livestock Science*, 119, 216–220.
- Hussain Z., Khan M.Z., Khan A., Javed I., Saleemi M.K., Mahmood S., Asi M.R. (2010): Residues of aflatoxin B1 in broiler meat: effect of age and dietary aflatoxin B1 levels. *Food and Chemical Toxicology*, 48, 3304–3307.
- Hussein H.S., Brasel J.M. (2001): Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*, 167, 101–134.
- Kermanshahi H., Akbari M., Maleki M., Behgar M. (2007): Effect of prolonged low level inclusion of aflatoxin B1 into diet on performance, nutrient digestibility, histopathology and blood enzymes of broiler chickens. *Journal of Animal and Veterinary Advances*, 6, 686–692.
- Magnoli A.P., Monge M.P., Miazzi R.D., Cavaglieri L.R., Magnoli C.E., Merck C.I., Cristofolini A.L., Dalcero A.M., Chiacchiera S.M. (2011): Effect of low levels of aflatoxin B on performance, biochemical parameters, and aflatoxin B in broiler liver tissues in the presence of monensin and sodium bentonite. *Poultry Science*, 90, 48–58.
- Marin D.E., Taranu I., Bunaciu R.P., Pascale F., Tudor D.S., Avram N., Sarca M., Cureu I., Criste R.D., Suta V. (2002): Changes in performance, blood parameters, humoral and cellular immune responses in weanling piglets exposed to low doses of aflatoxin. *Journal of Animal Science*, 80, 1250.
- Martinez-de-Anda A., Valdivia A.G., Jaramillo-Juarez F., Reyes J.L., Ortiz R., Quezada T., de Luna M.C., Rodriguez M.L. (2010): Effects of aflatoxin chronic intoxication in renal function of laying hens. *Poultry Science*, 89, 1622–1628.
- Meucci V., Razzuoli E., Soldani G., Massart F. (2010): Mycotoxin detection in infant formula milks in Italy. *Food Additives & Contaminants. Part A – Chemistry, Analysis, Control, Exposure & Risk Assessment*, 27, 64–71.
- Micco C., Miraglia M., Onori R., Brera C., Mantovani A., Ioppolo A., Stasolia D. (1988): Long-term administration of low doses of mycotoxins to poultry. 1. Residues of aflatoxin B1 and its metabolites in broilers and laying hens. *Food Additives and Contaminants*, 5, 303–308.
- Monbaliu S., Van Poucke C., Detavernier C., Dumoulin F., Van De Velde M., Schoeters E., Van Dyck S., Averkieva O., Van Peteghem C., De Saeger S. (2010): Occurrence of mycotoxins in feed as analyzed by a multi-mycotoxin LC-MS/MS method. *Journal of Agricultural and Food Chemistry*, 58, 66–71.
- National Research Council (1994): *Nutrient Requirements of Poultry*. 9<sup>th</sup> Ed. National Academy Press, Washington, USA.
- Nomura H., Ogiso M., Yamashita M., Takaku H., Kimura A., Chikaso M., Nakamura Y., Fujii S., Watai M., Yamada H. (2011): Uptake by dietary exposure and elimination of aflatoxins in muscle and liver of rainbow trout (*Oncorhynchus mykiss*). *Journal of Agricultural and Food Chemistry*, 59, 5150–5158.
- Oguz H., Kurtoglu V. (2000): Effect of clinoptilolite on performance of broiler chickens during experimental aflatoxicosis. *British Poultry Science*, 41, 512–517.
- Oliveira C.A.F., Kobashigawa E., Reis T.A., Mestieri L., Albuquerque R., Correa L.M.B. (2000): Aflatoxin B1 residues in eggs of laying hens fed a diet containing different levels of the mycotoxin. *Food Additives & Contaminants*, 17, 459–462.
- Oliveira C.A.F., Rosmaninho J.F., Castro A.L., Butkeraitis P., Reis T.A., Correa B. (2003): Aflatoxin residues in eggs of laying Japanese quail after long-term administration of rations containing low levels of aflatoxin B1. *Food Additives & Contaminants*, 20, 648–653.
- Ortatatli M., Oguz H., Karaman M. (2005): Evaluation of pathological changes in broilers during chronic aflatoxin (50 and 100 ppb) and clinoptilolite exposure. *Research in Veterinary Science*, 78, 61–68.
- Placinta C.M., D’Mello J.P.F., Macdonald A.M.C. (1999): A review of worldwide contamination of cereal grains

- and animal feed with *Fusarium* mycotoxins. *Animal Feed Science and Technology*, 78, 21–37.
- Reddy K.R., Reddy C.S., Muralidharan K. (2009): Detection of *Aspergillus* spp. and aflatoxin B1 in rice in India. *Food Microbiology*, 26, 27–31.
- Swenson D.H., Lin J.K., Miller E.C., Miller J.A. (1977): Aflatoxin B1-2, 3-oxide as a probable intermediate in the covalent binding of aflatoxins B1 and B2 to rat liver DNA and ribosomal RNA *in vivo*. *Cancer Research*, 37, 172.
- Tavcar-Kalcher G., Vrtac K., Pestevsek U., Vengust A. (2007): Validation of the procedure for the determination of aflatoxin B1 in animal liver using immunoaffinity columns and liquid chromatography with postcolumn derivatisation and fluorescence detection. *Food Control*, 18, 333–337.
- Van Pamel E., Verbeken A., Vlaemynck G., De Boever J., Daeseleire E. (2011): Ultrahigh-performance liquid chromatographic-tandem mass spectrometric multi-mycotoxin method for quantitating 26 mycotoxins in maize silage. *Journal of Agricultural and Food Chemistry*, 59, 9747–9755.
- Wogan G.N., Edwards G.S., Newberne P.M. (1971): Structure-activity relationships in toxicity and carcinogenicity of aflatoxins and analogs. *Cancer Research*, 31, 1936–1942.
- Wolzak A., Pearson A.M., Coleman T.H., Pestka J.J., Gray J.I. (1985): Aflatoxin deposition and clearance in the eggs of laying hens. *Food and Chemical Toxicology*, 23, 1057–1061.
- Wu F., Munkvold G.P. (2008): Mycotoxins in ethanol co-products: modeling economic impacts on the livestock industry and management strategies. *Journal of Agricultural and Food Chemistry*, 56, 3900–3911.
- Yang L., Ding H., Gu Z., Zhao J., Chen H., Tian F., Chen Y.Q., Zhang H., Chen W. (2009): Selection of single chain fragment variables with direct coating of aflatoxin B1 to enzyme-linked immunosorbent assay plates. *Journal of Agricultural and Food Chemistry*, 57, 8927–8932.
- Yunus A.W., Razzazi-Fazeli E., Bohm J. (2011): Aflatoxin B1 in affecting broiler's performance, immunity, and gastrointestinal tract: A review of history and contemporary issues. *Toxins*, 3, 566–590.
- Zhang Y., Caupert J. (2012): Survey of mycotoxins in U.S. distiller's dried grains with solubles from 2009 to 2011. *Journal of Agricultural and Food Chemistry*, 60, 539–543.
- Zhao J., Shirley R.B., Dibner J.D., Uraizee F., Officer M., Kitchell M., Vazquez-Anon M., Knight C.D. (2010): Comparison of hydrated sodium calcium aluminosilicate and yeast cell wall on counteracting aflatoxicosis in broiler chicks. *Poultry Science*, 89, 2147–2156.

Received: 2012–02–17

Accepted after corrections: 2012–06–07

---

*Corresponding Author*

Prof. Keying Zhang, Institute of Animal Nutrition, Sichuan Agricultural University, Yaan, Sichuan 625014, P.R.China  
Tel. +86 08 352 885 630, fax +86 08 352 885 630, e-mail: zkeying@yahoo.com

---